


ASSOCIATIONS BETWEEN SERUM LEVELS OF C3, C4, AND TOTAL CLASSICAL COMPLEMENT ACTIVITY IN COVID-19 PATIENTS AT THE TIME OF ADMISSION AND CLINICAL OUTCOME



A. Razi, A. Azimian, R. Arezumand, A. Solati, H.N. Ahmadabad

North Khorasan University of Medical Sciences, Bojnurd, Iran

Abstract. In the present study, we investigated the association between complement system status at the time of admission and clinical outcomes in COVID-19 patients. This single-center study was carried out with sixty-one adult patients with COVID-19 who were hospitalized at Imam Hassan Hospital of North Khorasan University of Medical Sciences (Bojnurd, Iran) with less than three days passage since onset of COVID-19 symptoms. Twenty-three healthy volunteers with demographic features similar to the patient group (matched by age and gender) were included in the study as a control group. Patient information including demographic information, demographic data, clinical characteristics, and clinical outcomes were obtained from electronic medical records. Of 61 hospitalized patients with COVID-19, 28 (47.54%) were female, and the average age was 48.7 ± 8.8 years. The healthy control group included 23 cases (11 (47.8%) female, 12 (52.1%) males, mean age 46.4 ± 4.4 years). Twenty-one of the 61 patients (34.4%) were admitted to the ICU, and sixteen of them (26.2%) died. Thirty-three (54.10%) patients with COVID-19 were hospitalized for less than 7 days, and 28 (45.90%) of them were hospitalized for ≥ 7 days. Our results show that length of hospital stay in the no-ICU group was significantly lower than the ICU admission or death groups (6.49 ± 0.24 vs. 8.85 ± 1.59 and 10.53 ± 1.80 , $p = 0.0002$). The levels of C3, C4, and CH50 were determined through the immunoturbidimetric method and single-radial-haemolysis plates, respectively, on serum samples obtained from patients at the time of admission or those in the control group. Our results indicate that C3, C4 and CH50 levels were markedly lower in COVID-19 patients than in the control group. We also found that complement parameter levels in COVID-19 patients who died or were admitted to ICU were significantly lower than in non-ICU COVID-19 patients. In general, it seems that serum level of C3, C4, and CH50 at admission may predict disease progression or adverse clinical outcome in COVID-19 patients.

Key words: COVID-19, complement system, clinical outcome, intensive care unit, discharge, mortality.

СВЯЗЬ УРОВНЯ СЫВОРОТОЧНЫХ С3, С4 И ОБЩЕЙ АКТИВНОСТИ КЛАССИЧЕСКОГО ПУТИ АКТИВАЦИИ КОМПЛЕМЕНТА У ПАЦИЕНТОВ С COVID-19 ПРИ ГОСПИТАЛИЗАЦИИ С КЛИНИЧЕСКИМ ИСХОДОМ

Рази А., Азимян А., Арезуманд Р., Солати А., Ахмадабад Х.Н.

Северо-Хорасанский университет медицинских наук, г. Боджнурд, Иран

Резюме. В статье представлены результаты оценки зависимости клинического исхода у пациентов с COVID-19 от состояния системы комплемента на момент госпитализации. В одноцентровое исследование были

Адрес для переписки:

Хасан Намдар Ахмадабад
Иран, г. Боджнурд, Медицинский университет Северного Хорасана.
Тел.: +98-5831513047. Факс: +98-5831513001
E-mail: namdar360@gmail.com

Contacts:

Hasan Namdar Ahmadabad
Iran, Bojnurd, North Khorasan University of Medical Sciences.
Tel.: +98-5831513047. Fax: +98-5831513001.
E-mail: namdar360@gmail.com

Для цитирования:

Рази А., Азимян А., Арезуманд Р., Солати А., Ахмадабад Х.Н. Связь уровня сывороточных С3, С4 и общей активности классического пути активации комплемента у пациентов с COVID-19 при госпитализации с клиническим исходом // Инфекция и иммунитет. 2022. Т. 12, № 5. С. 869–874. doi: 10.15789/2220-7619-ABS-1925

Citation:

Razi A., Azimian A., Arezumand R., Solati A., Ahmadabad H.N. Associations between serum levels of C3, C4, and total classical complement activity in COVID-19 patients at the time of admission and clinical outcome // Russian Journal of Infection and Immunity = Infektsiya i immunitet, 2022, vol. 12, no. 5, pp. 869–874. doi: 10.15789/2220-7619-ABS-1925

This work was supported by the North Khorasan University of Medical Sciences, Bojnurd, Iran (grant No. 1399/P/990001).

включены госпитализированные в больницу имама Хасана Северо-Хорасанского университета медицинских наук (Боджнурд, Иран) взрослые пациенты с COVID-19 ($n = 61$), у которых с момента появления симптомов COVID-19 прошло менее трех дней. Контрольную группу составили 23 здоровых добровольца, сходных по возрасту и полу с основной группой. Информация о пациентах, включая данные о возрасте, половой принадлежности, особенностях клинического течения и исходах заболевания, была получена из электронных медицинских карт. Из числа госпитализированных с COVID-19 пациентов 47,54% составляли женщины (28 из 61), средний возраст пациентов — $48,7 \pm 8,8$ года. В контрольную группу вошли 23 человека (11 (47,8%) женщин и 12 (52,1%) мужчин), средний возраст — $46,4 \pm 4,4$ года. Двадцать один из 61 пациента (34,4%) был направлен в отделение интенсивной терапии (ОИТ), из них шестнадцать (26,2%) умерли. Продолжительность госпитализации 33 (54,1%) пациентов с COVID-19 составила < 7 дней, 28 (45,9%) пациентов находились в стационаре ≥ 7 дней. Показано, что продолжительность пребывания в стационаре в группе пациентов, не нуждавшихся в оказании медицинской помощи в ОИТ, была значительно ниже, чем в группах пациентов, помещенных в ОИТ, или умерших ($6,49 \pm 0,24$ против $8,85 \pm 1,59$ и $10,53 \pm 1,8$ соответственно, $p = 0,0002$). Уровни C3, C4 определяли иммунотурбидиметрическим методом, уровень CH50 — одномерным радиальным гемолизом. Материалом для исследований служили образцы сыворотки, полученные от пациентов при поступлении в стационар и от лиц контрольной группы. Установлено, что у пациентов с COVID-19 уровни C3, C4 и CH50 были существенно ниже, чем у лиц контрольной группы, а у пациентов ОИТ и умерших — ниже, чем у пациентов с COVID-19, не нуждавшихся в терапии в условиях ОИТ. Таким образом, возможно, что уровень сывороточных C3, C4 и CH50 при поступлении может являться прогностическим фактором дальнейшего течения заболевания и развития неблагоприятных клинических исходов у пациентов с COVID-19.

Ключевые слова: COVID-19, система комплемента, клинический результат, отделение интенсивной терапии, выписка, смертность.

Introduction

More than a year after its emergence, COVID-19, the disease caused by SARS-CoV-2, continues to plague the world and dominate our daily lives [2]. Patients with COVID-19 may develop symptoms such as fever, dry cough, pharyngeal pain, tiredness, abdominal pain, diarrhea, conjunctivitis, muscle fatigue, or pneumonia; some are left with serious side effects or even die [14]. Previous studies have reported that 14.1–33.0% of COVID-19 patients are prone to develop into severe cases, and the mortality rate of critical cases is 61.5%, increasing sharply with age and underlying comorbidities [22, 23]. In more severe cases of COVID-19, patients can develop acute respiratory distress syndrome (ARDS), leading to a worse prognosis [9]. Deregulated activation of multiple adaptive and innate immune pathways (including T cell activation, cytokine expression from macrophages and neutrophils, the complement system, and several pro-coagulant and thrombogenic pathways) is believed to fuel a hyperinflammatory state that drives ARDS and may lead to multiple organ injury and finally death in COVID-19 [13, 19].

The complement system is a key part of the innate immune system which plays an important role in defense against foreign pathogens such as viruses but, in addition to being an important part of the immune defense system, it plays a critical role in promoting the inflammatory process that leads to organ dysfunction [21]. Although several studies have been carried out on the complement system in COVID-19 and its relationship with clinical outcomes [5, 6, 8,

12, 25], they did not indicate a clear protective or adverse effect of this system. Dheiret al. in a retrospective study showed there is no significant difference, in terms of C3 and C4 levels, in both ICU and non-ICU COVID-19 patients [5]. They suggested that measurement of C3 and C4 levels cannot be used to show severity disease [5].

Controversially, Ghazavi et al. found that the levels of C3 and C4 in non-severe COVID-19 patients were significantly higher than in severe-COVID-19 patients [8]. A previous study by Fang et al. also indicated decreased complement C3 levels are associated with poor prognosis in COVID-19 patients [6]. Java and colleagues claimed that the role of the complement system in COVID-19 patients is time dependent, wherein complement activation in the first week of infection can serve as a “friend”, and its activation in the second or third weeks of infection can be a “foe” [12]. Zinellu et al., in a systematic review, concluded that lower concentrations of C3 and C4 are significantly associated with higher COVID-19 severity and mortality [25]. They suggested that additional studies are required to determine whether measurement of complement components can be useful to predict adverse clinical consequences in COVID-19 patients [25].

Therefore, we conducted a study to assess the association between complement system status at the time of admission and clinical outcomes (e.g., length of stay, ICU admission, discharge, mortality) in COVID-19 patients. Understanding this association will help us elucidate the role of the complement system for prediction of the risk of developing critical COVID-19.

Materials and methods

Study design and participants. This single-center study was carried out with sixty-one adult patients with COVID-19 who were hospitalized at Imam Hassan Hospital of North Khorasan University of medical sciences (Bojnourd, Iran) from April 20 to August 5, 2021 with less than three days passage since the onset of COVID-19 symptoms. All patients with a positive RT-PCR test for SARS-CoV-2 and common CT imaging findings associated with COVID-19 were included in the study. Patients also had no history of SARS-CoV-2 infection or COVID-19 vaccination.

Available data suggest that patients with mild-to-moderate COVID-19 remain infectious no longer than three days after symptom onset. Recent evidence indicates that age and comorbidities could possibly confound the association between complement system status and clinical events [16]. Therefore, we adjusted for sex, age and comorbidities. Twenty-three healthy volunteers with demographic features similar to the patient group (matched by age and gender) were included in the study as a control group.

Data collection and assessment of serum complement levels and activity. We obtained demographic data, exposure history, chronic medical histories, clinical symptoms or signs, clinical outcomes, and hospitalization duration from electronic medical records. We also collected serum samples from PCR-confirmed COVID-19 patient samples sent to Imam Hasan Hospital laboratory on the first day of admission for analysis of serum levels of C3, C4, and total classical complement activity (CH50 assay) present in the serum. We also collected blood samples from the healthy controls with a negative PCR-test for COVID-19. Blood samples were centrifuged immediately, and serum was obtained and frozen at -70°C until use.

Serum C3 and C4 concentrations were determined using kits from Roche Diagnostics (Indianapolis, IN, USA) according to the immunoturbidimetric method [18]. The CH50 test was evaluated using single-radial-haemolysis plates (Biogen, Iran) as described elsewhere [18]. All values were compared to the normal ranges which were reported as: 89–187 mg/dL for C3; 10–40 mg/dL for C4; and 70–130 mm for CH50.

Statistical analysis. All statistical analyses were carried out using GraphPad Prism 5.0 (GraphPad, San Diego, CA, USA). Data distribution was analyzed by a Kolmogorov-Smirnov test. According to the results of the normality test, a one-way ANOVA followed by Dunn's or Tukey's post-hoc test, or a non-parametric Kruskal-Wallis test, were used for statistical comparisons. Analyses were adjusted for age, sex and comorbidities. Data were expressed as mean \pm standard deviation (SD). Values of $p < 0.05$ (*) were considered significant.

Results

Demographic data and clinical characteristics. Of 61 hospitalized patients with COVID-19, 28 (47.54%) were female, and the average age was 48.7 ± 8.8 years. The most commonly self-reported symptoms at onset of illness were fever ($n = 41$ [67.2%]), cough ($n = 39$ [63.9%]), fatigue or myalgia ($n = 19$ [31.4%]), diarrhea ($n = 9$ [14.7%]), or headache ($n = 6$ [9.8%]). Twenty-six (42.6%) patients had comorbidities, including cardiovascular disease ($n = 10$ [38.4%]), diabetes ($n = 6$ [23.0%]), hypertension ($n = 7$ [26.9%]), chronic kidney disease ($n = 3$ [11.5%]), and endocrine system diseases ($n = 2$ [7.6%]). The healthy control group included 23 cases (11 (47.8%) female, 12 (52.1%) males, mean age 46.4 ± 4.4 years).

Twenty-one of the 61 patients (34.4%) were admitted to the ICU, and sixteen of them (26.2%) died. Thirty-three (54.10%) patients with COVID-19 were hospitalized for less than 7 days, and 28 (45.90%) of them were hospitalized for ≥ 7 days. Our results show that length of hospital stay in the non-ICU group was significantly lower than in the ICU admission or death groups (6.49 ± 0.24 vs. 8.85 ± 1.59 and 10.53 ± 1.80 , $p = 0.0002$).

C3, C4, and CH50 levels in COVID-19 patients and the healthy control group. As shown in Table, serum C3 and C4 concentrations and CH50 assay were markedly lower in COVID-19 patients than in the healthy control group. Further statistical analysis showed that serum levels of C3, C4, and CH50 in the non-ICU admission group were statistically higher than in the death and ICU admission groups. We did not observe a significant difference in C3, C4, and CH50 levels between ICU admission and death groups of COVID-19 patients ($p > 0.05$).

We also analyzed the relationship between C3, C4, and CH50 levels in COVID-19 patients and hospital length of stay. As shown in Figure, serum levels of C3, C4, and CH50 at the time of admission in COVID-19 patients who were hospitalized for 7 or more days were statistically lower than in COVID-19 patients who were hospitalized less than 7 days ($p < 0.05$).

Discussion

The complement system efficiently recognizes and eliminates viral pathogens via several mechanisms including: opsonization of viruses; lysing of virus-infected cells; induction of an antiviral immunoinflammatory state; boosting of virus-specific immune responses; and directly neutralizing cell-free viruses [1]. On the other hand, complement activation in viral infections may play a critical role in pathogenesis, clinical manifestation, and disease severity [3, 10].

Several reports have shown that the concentration of complement components, and serum complement

Table. Comparison of serum C3, C4 and CH50 levels at the time of admission between COVID-19 patients with different clinical outcomes and the healthy control group

Lab Variable	Healthy Control	No-ICU Admission	ICU Admission	Death	P value
C3, mg/dl	137.1±25.82	98.76±19.09	72.62±13.92	71.63±14.28	< 0.0001
C4, mg/dl	28.50±6.25	24.46±5.87	17.71±6.96	18.79±5.44	< 0.0001
CH50, mm	94.43±13.66	91.58±17.89	77.24±12.66	78.50±7.64	0.007


activity, change in the COVID-19 patients. However, they did not definitively determine whether measurement of complement components or serum complement activity can be useful to predict adverse clinical outcomes in COVID-19 patients. Therefore, this study was designed to assess the relationship between serum levels of C3, C4, and total classical complement activity in COVID-19 patients at the time of admission and clinical outcomes. The results of this study indicate that serum levels of C3 and C4 and CH50 assay were markedly lower in COVID-19 patients than in the healthy control group. The most interesting finding was that serum levels of C3, C4, and total classical complement activity in COVID-19 patients who died or were ICU admitted were significantly lower than in COVID-19 patients who were not ICU admitted.

Conversely, Keshavarz and colleagues showed that serum levels of the C3 and C4 factors have no significant change between patients and healthy individuals [15]. In another study, Zhang et al. showed that complement C3 cannot predict disease progression [24]. Our findings are also contrary to that of Dheirs et al. who found there was no significant difference in terms of C3 and C4 levels in both ICU and non-ICU COVID-19 patients [5]. Henry et al. have also shown that complement hyperactivation failed to predict progression to severe COVID-19 [11]. In addition, they also showed that there are no significant differences in total classical complement activity (or CH50 level) at the time of admission between COVID-19 patients with different clinical outcomes [11]. These inconsistencies

may be due to differences in study design, limited sample size, characteristics of COVID-19 patients, time between the onset of symptoms and obtaining the blood sample, or confounders.

However, this study supports the findings from previous observations [6, 8, 15]. Fang et al. in a retrospective cohort study investigated C3 levels in COVID-19 patients [6]. They showed that C3 levels in the non-survival group were significantly lower than in the survival group; they concluded that decreased complement C3 levels are associated with poor prognosis in COVID-19 patients [6]. The finding also match earlier findings by Ghazavi et al., which showed that C3 and C4 levels were markedly decreased in severe COVID-19 patients in comparison with non-severe COVID-19 patients [8]. In agreement with our data, Keshavarz et al. indicated that mean CH50 activity level in COVID-19 patients is significantly reduced compared to healthy individuals [15].

The decreased levels of C3 and C4 in COVID-19 patients (in comparison to the healthy group), and significant differences between the non-ICU admission group and the death and ICU admission groups, may be explained by the fact that hyperactivation of the complement system leads to proteolytic cleavage of the key complement molecules C3 and C4, leading to cleavage products including C3a, C3b, C4a, and C4b. These may trigger severe inflammatory responses in numerous organs [4]. In support of this possibility, Brandon and colleagues have demonstrated that C3a and the C3a/C3 ratio are significantly elevated in severe COVID-19 patients

**Figure. Comparison of hospital stay length in COVID-19 patients with serum levels of C3, C4 and CH50 at the time of admission**

Note. Data are mean±SD. *p < 0.05 was considered statistically significant.

presenting to the emergency department compared to mild or moderate severity COVID-19 patients [11]. On the other hand, Fletcher-Sandersjö et al. suggested that overactivation of the complement cascade in COVID-19 patients is associated with activation of coagulation systems and consequent severe complications [7]. It is possible, therefore, that overproduction of C3a and C4a following complement hyperactivation is associated with adverse outcomes in patients with SARS-CoV-2 infection through activation of the coagulation cascade and thrombus formation.

Taken together, these results suggest that the levels of complement C3, C4, and CH50 at admission may predict disease progression and adverse clinical outcome in patients with SARS-CoV-2 infection. These findings may be somewhat limited by several confounders, such as different viral strains, sociodemographic factors, and various types of drug therapy during hospitalization [17, 20]. Thus, we suggest that future studies include: investigation of alteration of C3 and C4 at different stages of the disease; determination of the relationship between serum levels of C3 and C4 and inflammatory biomarkers; and design of clinical trials with complement inhibitors, such as a C3 inhibitor.

References

1. Agrawal P., Nawadkar R., Ojha H., Kumar J., Sahu A. Complement evasion strategies of viruses: an overview. *Front. Microbiol.*, 2017, vol. 8, pp. 1117. doi: 10.3389/fmicb.2017.01117
2. Baric R.S. Emergence of a highly fit SARS-CoV-2 variant. *N. Engl. J. Med.*, 2020, vol. 383, no. 27, pp. 2684–2686. doi: 10.1056/NEJMcibr2032888
3. Bjornson A.B., Mellencamp M.A., Schiff G.M. Complement is activated in the upper respiratory tract during influenza virus infection. *Am. Rev. Respir. Dis.*, 1991, vol. 143, no. 5, pp. 1062–1066. doi: 10.1164/ajrccm/143.5_Pt_1.1062
4. Bosmann M., Ward P.A. Role of C3, C5 and anaphylatoxin receptors in acute lung injury and in sepsis. In: Current Topics in Innate Immunity I.I. Advances in Experimental Medicine and Biology, vol. 946. Eds.: J. Lambris, G. Hajishengallis. *New York: Springer*, 2012, vol. 946, pp. 147–159. doi: 10.1007/978-1-4614-0106-3_9
5. Dheir H., Sipahi S., Yaylaci S., Koroğlu M., Erdem A.F., Karabay O. Is there relationship between SARS-CoV-2 and the complement C3 and C4? *Turk. J. Med. Sci.*, 2020, vol. 50, no. 4, pp. 687–688. doi: 10.3906/sag-2004-336
6. Fang S., Wang H., Lu L., Jia Y., Xia Z. Decreased complement C3 levels are associated with poor prognosis in patients with COVID-19: a retrospective cohort study. *Int. Immunopharmacol.*, 2020, vol. 89 (Pt A): 10707. doi: 10.1016/j.intimp.2020.10707
7. Fletcher-Sandersjö A., Bellander B.-M. Is COVID-19 associated thrombosis caused by overactivation of the complement cascade? A literature review. *Thromb. Res.*, 2020, vol. 194, pp. 36–41. doi: 10.1016/j.thromres.2020.06.027
8. Ghazavi A., Mosayebi G., Keshavarzian N., Rabiemajid S., Ganji A. Reduction of inflammatory C3 and C4 complement proteins in severe COVID-19 patients. *Preprint (Version 1) available at Research Square*. 2020. 13 p. doi: 10.21203/rs.3.rs-127493/v1
9. Gibson P.G., Qin L., Puah S.H. COVID-19 acute respiratory distress syndrome (ARDS): clinical features and differences from typical pre-COVID-19 ARDS. *Med. J. Aust.*, 2020, vol. 213, no. 2, pp. 54–56. doi: 10.5694/mja2.50674
10. Gralinski L.E., Sheahan T.P., Morrison T.E., Menachery V.D., Jensen K., Leist S.R., Whitmore A., Heise M.T., Baric R.S. Complement activation contributes to severe acute respiratory syndrome coronavirus pathogenesis. *mBio*, 2018, vol. 9, no. 5: e01753-18 doi: 10.1128/mBio.01753-18
11. Henry B.M., Szerguyk I., de Oliveira M.H.S., Lippi G., Benoit J.L., Vikse J., Benoit S.W. Complement levels at admission as a reflection of Coronavirus Disease 19 (COVID-19) severity state. *J. Med. Virol.*, 2021, vol. 93, no. 9, pp. 5515–5522. doi: 10.1002/jmv.27077
12. Java A., Apicelli A.J., Liszewski M.K., Coler-Reilly A., Atkinson J.P., Kim A.H., Kulkarni H.S. The complement system in COVID-19: friend and foe? *JCI Insight*, 2020, vol. 5, no. 15: e140711. doi: 10.1172/jci.insight.140711
13. Jayarangaiah A., Kariyanna P.T., Chen X., Jayarangaiah A., Kumar A. COVID-19-associated coagulopathy: an exacerbated immunothrombosis response. *Clin. Appl. Thromb. Hemost.*, 2020, vol. 26: 1076029620943293. doi: 10.1177/1076029620943293
14. Jimenez-Cebrian A.M., Castro-Mendez A., García-Podadera B., Romero-Galisteo R., Medina-Alcántara M., Garcia-Paya I., Páez-Moguer J., Córdoba-Fernández A. Clinical manifestations of COVID-19 in the feet: a review of reviews. *J. Clin. Med.*, 2021, vol. 10, no. 10: 2201. doi: 10.3390/jcm10102201
15. Keshavarz F., Ghalamfarsa F., Javdansirat S., Hasanzadeh S., Azizi A., Sabz G., Salehi M., Ghalamfarsa G. Patients with COVID-19 have significantly reduced CH50 activity. *Virusdisease*, 2021, vol. 32, no. 4, pp. 681–689. doi: 10.1007/s13337-021-00710-6

Acknowledgments

This work was supported by the North Khorasan University of Medical Sciences, Bojnurd, Iran. The authors thank the members of the Department of Pathobiology and Medical Laboratory Sciences, North Khorasan University of Medical Sciences, Bojnurd, Iran for their technical assistance. We would also like to thank the clinical staff of Imam Hassan Hospital for assisting us in this research project.

Ethics approval

The study was performed based on an informed consent and was approved by the Ethics Committee of North Khorasan University of Medical Sciences, Bojnourd, Iran (IR.NKUMS.REC.1399.021).

Informed consent

Informed consent was obtained from all individual participants included in the study.

Conflict of interest

The authors declare that they have no conflict of interest.

16. Kristensen M.K., Hansen M.B., Madsen M.B., Hansen C.B., Pilely K., Hyldegaard O., Garred P. Complement activation is associated with mortality in patients with necrotizing soft-tissue infections — a prospective observational study. *Front. Immunol.*, 2020, vol. 11: 17. doi: 10.3389/fimmu.2020.00017
17. Lippi G., Henry B.M., Hoehn J., Benoit S., Benoit J. Validation of the Corona-Score for rapid identification of SARS-CoV-2 infections in patients seeking emergency department care in the United States. *Clin. Chem. Lab. Med.*, 2020, vol. 8, no. 12, pp. e311–e113. doi: 10.1515/cclm-2020-1121
18. McPherson R.A., Pincus M.R. Henry's clinical diagnosis and management by laboratory methods. *Elsevier Health Sciences*, 2021. 1618 p. URL: <https://www.elsevier.com/books/henrys-clinical-diagnosis-and-management-by-laboratory-methods/mcpherson/978-0-323-67320-4> (10.02.2022)
19. Melenotte C., Silvin A., Goubet A.G., Lahmar I., Dubuisson A., Zumla A., Raoult D., Merad M., Gachot B., Hénon C., Solary E., Fontenay M., André F., Maeurer M., Ippolito G., Piacentini M., Wang F.S., Ginhoux F., Marabelle A., Kroemer G., Derosa L., Zitvogel L. Immune responses during COVID-19 infection. *Oncoimmunology*, 2020, vol. 9, no. 1: 1807836. doi: 10.1080/2162402X.2020.1807836
20. Qureshi A.I., Baskett W.I., Huang W., Shyu D., Myers D., Lobanova I., Naqvi S.H., Thompson V.S., Shyu C.R. Effect of race and ethnicity on in-hospital mortality in patients with COVID-2019. *Ethn. Dis.*, 2021, vol. 31, no. 3, pp. 389–398. doi: 10.18865/ed.31.3.389
21. Raghunandan S., Josephson C.D., Verkerke H., Linam W.M., Ingram T.C., Zerra P.E., Arthur C.M., Stowell S.R., Briones M., Chonat S. Complement inhibition in severe COVID-19 acute respiratory distress syndrome. *Front. Pediatr.*, 2020, vol. 8: 616731. doi: 10.3389/fped.2020.616731
22. Shakeri H., Azimian A., Ghasemzadeh-Moghaddam H., Safdari M., Haresabadi M., Daneshmand T., Namdar Ahmadabad H. Evaluation of the relationship between serum levels of zinc, vitamin B12, vitamin D, and clinical outcomes in patients with COVID-19. *J. Med. Virol.*, 2022, vol. 94, no. 1, pp. 141–146. doi: 10.1002/jmv.27277
23. Xu W., Sun N.N., Gao H.N., Chen Z.Y., Yang Y., Ju B., Tang L.L. Risk factors analysis of COVID-19 patients with ARDS and prediction based on machine learning. *Sci. Rep.*, 2021, vol. 11, no. 1: 2933. doi: 10.1038/s41598-021-82492-x
24. Zhang Z., Li X., Zhang W., Shi Z.-L., Zheng Z., Wang T. Clinical features and treatment of 2019-nCoV pneumonia patients in Wuhan: report of a couple cases. *Virol. Sin.*, 2020, vol. 35, no. 3, pp. 330–336. doi: 10.1007/s12250-020-00203-8
25. Zinellu A., Mangoni A.A. Serum complement C3 and C4 and COVID-19 severity and mortality: a systematic review and meta-analysis with meta-regression. *Front. Immunol.*, 2021, vol. 12: 696085. doi: 10.3389/fimmu.2021.696085

Авторы:

Рази А., д.м.н., доцент, преподаватель кафедры урологии, больница имама Али, Северо-Хорасанский университет медицинских наук, г. Боджнурд, Иран;

Азимян А., д.м.н., доцент, преподаватель кафедры патобиологии и лабораторных наук медицинского факультета Северо-Хорасанского университета медицинских наук, г. Боджнурд, Иран;

Арезуманд Р., д.м.н., ассистент, преподаватель кафедры патобиологии и лабораторных наук медицинского факультета Северо-Хорасанского университета медицинских наук, г. Боджнурд, Иран;

Солати А., д.ф.н., доцент, преподаватель кафедры английского языка медицинского факультета Северо-Хорасанского университета медицинских наук, г. Боджнурд, Иран;

Ахмадабад Х.Н., д.м.н., доцент, преподаватель кафедры патобиологии и лабораторных наук медицинского факультета Северо-Хорасанского университета медицинских наук, г. Боджнурд, Иран.

Authors:

Razi A., PhD, MD (Medicine), Assistant Professor, Faculty Member, Department of Urology, Imam Ali Hospital, North Khorasan University of Medical Sciences, Bojnurd, Iran;

Azimian A., PhD, MD (Medicine), Associate Professor, Faculty Member, Department of Pathobiology and Laboratory Sciences, School of Medicine, North Khorasan University of Medical Sciences, Bojnurd, Iran;

Arezumand R., PhD, MD (Medicine), Assistant Professor, Faculty Member, Department of Pathobiology and Laboratory Sciences, School of Medicine, North Khorasan University of Medical Sciences, Bojnurd, Iran;

Solati A., PhD, MD (Philosophy), Associate Professor, Faculty Member, Department of English Language, School of Medicine, North Khorasan University of Medical Sciences, Bojnurd, Iran;

Ahmadabad H.N., PhD, MD (Medicine), Associate Professor, Faculty Member, Department of Pathobiology and Laboratory Sciences, School of Medicine, North Khorasan University of Medical Sciences, Bojnurd, Iran.

Поступила в редакцию 08.04.2022
Отправлена на доработку 15.05.2022
Принята к печати 27.06.2022

Received 08.04.2022
Revision received 15.05.2022
Accepted 27.06.2022