

SEROPREVALENCE OF TICK-BORNE DISEASES IN THE NORTHWEST FEDERAL DISTRICT OF THE RUSSIAN FEDERATION



N.K. Tokarevich^a, O.V. Blinova^a, N.A. Stoyanova^a, R.R. Baimova^a, E.A. Siuziumova^a, V.I. Lomonosova^a, A.A. Tronin^b, R.V. Buzinov^c, O.V. Sokolova^c, B.R. Gnativ^d, L.V. Buts^e, L.A. Bubnova^f, O.S. Safonova^f, A.I. Stankevich^g, E.L. Kalinina^h, R. Vikseⁱ, A.K. Andreassenⁱ

^a St. Petersburg Pasteur Institute, St. Petersburg, Russian Federation

^b Scientific Research Centre for Ecological Safety of the Russian Academy of Sciences, St. Petersburg Federal Research Center of the Russian Academy of Sciences (SPC RAS), St. Petersburg, Russian Federation

^c Federal Service for Supervision of Consumer Rights Protection and Human Welfare in Arkhangelsk Oblast, Arkhangelsk, Russian Federation

^d Federal Service for Supervision of Consumer Rights Protection and Human Welfare in the Republic of Komi, Syktyvkar, Russian Federation

^e Federal Service for Supervision of Consumer Rights Protection and Human Welfare in Leningrad Oblast, St. Petersburg, Russian Federation

^f Center of Hygiene and Epidemiology in the Karelia Republic, Petrozavodsk, Russian Federation

^g Center of Hygiene and Epidemiology in Pskov Oblast, Pskov, Russian Federation

^h Federal Service for Supervision of Consumer Rights Protection and Human Welfare in Pskov Oblast, Pskov, Russian Federation

ⁱ Norwegian Institute of Public Health, Oslo, Norway

Abstract. *Introduction.* Knowledge about tick-borne disease (TBD) distribution is necessary to improve prevention, where-as detection of human serum IgG antibodies against relevant pathogens is a method for monitoring TBD prevalence in local population. The study objective was to estimate seroprevalence of IgG antibodies against tick-borne encephalitis virus (TBEV), *Borrelia burgdorferi sensu lato*, *Coxiella burnetii*, *Anaplasma phagocytophilum*, and *Ehrlichia chaffeensis/E. muris* in healthy residents from the five territories of the Northwestern Federal District of the Russian Federation (Arkhangelsk Oblast, Leningrad Oblast, Pskov Oblast, the Republic of Komi and the Republic of Karelia). *Materials and methods.* In 2017–2019, a total of 1244 serum samples from healthy residents, not vaccinated against TBDs or other flavivirus-caused infections was studied by ELISA. *Results.* 21.7% of the sera samples contained IgG antibodies against a single TBD pathogen, whereas 2.1% showed signs of coinfection with two or more pathogens. The most common were IgG antibodies against TBEV (5 territories, 12.2%), followed by *Borrelia burgdorferi sensu lato* (5 territories, 3.5%), *C. burnetii* (4 territories, 2.9%), *Anaplasma phagocytophilum* (3 territories, 1.6%), *E. chaffeensis/E. muris* (5 territories, 1.5%). The IgG antibodies were more common in men (55.2%) than in women (44.8%), being found virtually evenly in age-independent manner (from juniors under 18 to seniors over 60). *Conclusion.* The results of this first comprehensive serosurveillance study in the Northwestern Federal District of the Russian Federation assessing serum IgG antibodies against tick-borne diseases in-

Адрес для переписки:

Токаревич Николай Константинович
197101, Россия, Санкт-Петербург, ул. Мира, 14,
ФБУН НИИ эпидемиологии и микробиологии имени Пастера.
Тел.: 8 (812) 232-21-36.
E-mail: zoonoses@mail.ru

Contacts:

Nikolay K. Tokarevich
197101, Russian Federation, St. Petersburg, Mira str., 14,
St. Petersburg Pasteur Institute.
Phone: +7 (812) 232-21-36.
E-mail: zoonoses@mail.ru

Для цитирования:

Токаревич Н.К., Блинова О.В., Стоянова Н.А., Баимова Р.Р.,
Сюзюмова Е.А., Ломоносова В.И., Тронин А.А., Бузинов Р.В.,
Соколова О.В., Гнатив Б.Р., Бут Л.В., Бубнова Л.А., Сафонова О.С.,
Станкевич А.И., Калинина Е.Л., Викс Р., Андреассен А.К.
Серопревалентность клещевых заболеваний в Северо-Западном
федеральном округе Российской Федерации // Инфекция и иммунитет.
2022. Т. 12, № 5. С. 891–901. doi: 10.15789/2220-7619-SOT-1953

Citation:

Tokarevich N.K., Blinova O.V., Stoyanova N.A., Baimova R.R.,
Siuziumova E.A., Lomonosova V.I., Tronin A.A., Buzinov R.V., Sokolova O.V.,
Gnativ B.R., Buts L.V., Bubnova L.A., Safonova O.S., Stankevich A.I.,
Kalinina E.L., Vikse R., Andreassen A.K. Seroprevalence of tick-borne
diseases in the Northwest Federal District of the Russian Federation //
Russian Journal of Infection and Immunity = Infektsiya i immunitet, 2022,
vol. 12, no. 5, pp. 891–901. doi: 10.15789/2220-7619-SOT-1953

dicate a wide distribution of such pathogens. Moreover, infections caused by *C. burnetii*, *Anaplasma phagocytophilum*, and *Ehrlichia chaffeensis/E. muris* might be highly underdiagnosed.

Key words: seroprevalence, antibodies, tick-borne encephalitis, Lyme disease, human granulocytic anaplasmosis, human monocytic ehrlichiosis, Q fever, European North of Russia.

СЕРОПРЕВАЛЕНТНОСТЬ КЛЕЩЕВЫХ ЗАБОЛЕВАНИЙ В СЕВЕРО-ЗАПАДНОМ ФЕДЕРАЛЬНОМ ОКРУГЕ РОССИЙСКОЙ ФЕДЕРАЦИИ

Токаревич Н.К.¹, Блинова О.В.¹, Стоянова Н.А.¹, Баимова Р.Р.¹, Сюзюмова Е.А.¹, Ломоносова В.И.¹, Тронин А.А.², Бузинов Р.В.³, Соколова О.В.³, Гнатив Б.Р.⁴, Буц Л.В.⁵, Бубнова Л.А.⁶, Сафонова О.С.⁶, Станкевич А.И.⁷, Калинина Е.Л.⁸, Виксе Р.⁹, Андреассен А.К.⁹

¹ФБУН НИИ эпидемиологии и микробиологии имени Пастера, Санкт-Петербург, Россия

²Научно-исследовательский центр экологической безопасности Российской академии наук, Санкт-Петербургский федеральный исследовательский центр Российской академии наук (НИЦ РАН), Санкт-Петербург, Россия

³Федеральная служба по надзору в сфере защиты прав потребителей и благополучия человека по Архангельской области, г. Архангельск, Россия

⁴Федеральная служба по надзору в сфере защиты прав потребителей и благополучия человека по Республике Коми, г. Сыктывкар, Россия

⁵Федеральная служба по надзору в сфере защиты прав потребителей и благополучия человека по Ленинградской области, Санкт-Петербург, Россия

⁶ФБУЗ Центр гигиены и эпидемиологии Республики Карелия, г. Петрозаводск, Россия

⁷ФБУЗ Центр гигиены и эпидемиологии в Псковской области, г. Псков, Россия

⁸Федеральная служба по надзору в сфере защиты прав потребителей и благополучия человека по Псковской области, г. Псков, Россия

⁹Норвежский институт общественного здравоохранения, Осло, Норвегия

Резюме. Введение. Знание распространения клещевых заболеваний необходимо для улучшения их профилактики, а выявление IgG-антител к их возбудителям в сыворотках крови человека является методом мониторинга распространенности клещевых заболеваний среди местного населения. Цель исследования: изучение серопревалентности практически здорового населения Архангельской, Ленинградской, Псковской, областей, Республик Коми и Карелии, расположенных на европейском Севере России, в отношении вируса КЭ, *Borrelia burgdorferi sensu lato*, *Coxiella burnetii*, *Anaplasma phagocytophilum* и *Ehrlichia chaffeensis/E. muris*. Материалы и методы. Всего было исследовано 1244 сыворотки крови практически здоровых лиц, постоянно проживающих на данной территории, не вакцинированных против клещевого энцефалита и других инфекций, вызываемых флавивирусами, профессия которых не связана с риском заражения клещевыми инфекциями. Сыворотки крови доноров исследовали методом ELISA. Результаты. Установлено, что в 21,7% образцах исследованных сывороток, полученных от жителей всех анализируемых территорий, обнаружены антитела класса IgG к возбудителям различных инфекций, передающихся клещами, в том числе в 26 пробах — одновременно к двум и более патогенам. В сыворотках жителей республик Коми и Карелия, а также Ленинградской области выявлены антитела ко всем исследованным антигенам, в сыворотках жителей Архангельской области — антитела к вирусу клещевого энцефалита, *Borrelia burgdorferi sensu lato*, *C. urretiid* и *E. chaffeensis/E. muris* а в сыворотках жителей Псковской области — антитела к вирусу клещевого энцефалита, *Borrelia burgdorferi sensu lato* и *E. chaffeensis/E. muris*. У мужчин IgG-антитела к клещевым инфекциям определялись чаще (55,2%), чем у женщин (44,8%). IgG-антитела к клещевым инфекциям были определены практически равномерно у жителей всех возрастов от детского (до 18 лет) до пожилого (старше 60 лет). Обсуждение. Представленные результаты первого комплексного обследования жителей европейского Севера России на наличие в их сыворотках крови IgG-антител к клещевым инфекциям, свидетельствуют о широком распространении этих инфекций и, вероятно, о существенной гиподиагностике инфекций, вызываемых *C. urretiid*, *Anaplasma phagocytophilum* и *Ehrlichia chaffeensis/E. muris*.

Ключевые слова: серопревалентность, антитела, клещевой энцефалит, болезнь Лайма, гранулоцитарный анаплазмоз человека, моноцитарный эрлихиоз человека, Ку-лихорадка, Европейский Север России.

Introduction

TBDs are among important drivers of habitat quality, public and herd health, as well as the epidemiological safety of the local population [5]. Recent social changes and stronger anthropogenic impacts on abiotic components of landscapes may contrib-

ute to the evolution of those infections [1, 24]. In this regard, any advance in understanding of TBDs real distribution within a territory is of great importance for prevention of these diseases.

TBD incidence rate is an important factor that defines prophylaxis volume and methods within an administrative unit. However, it does not always re-

flect to a full degree the real spread of TBD, since that depends much on the awareness of the local population, their health seeking behavior, availability and quality of medical care, qualification of medical personnel, availability of laboratory diagnostics, etc. There is a high risk of underdiagnosis since most of the human TBDs appear as subclinical forms [5]. Many patients are asymptomatic or experience mild symptoms and, as a rule, do not seek medical help, thus reducing the reported incidence rate. However, the absence of pronounced symptoms does not exclude the probability of chronic TBD course, resulting in disability or even fatal outcome [1, 8, 24, 48]. Knowledge of TBDs actual distribution is necessary to improve its effective prevention in the area. The detection of IgG antibodies to TBD pathogens in blood sera of humans is a method for indicating the real level of infection in the local population.

Among all TBDs it is TBE that poses the most socially important problem in the Russian Federation, being diagnosed in 46 subdivisions. At least 1508 cases of TBE were reported in 2018, including 153 in children under 14. TBE incidence rate was 1.3 per 100 thousand inhabitants; in 98% of cases the disease developed after tick bites. Twenty two cases of TBE were fatal; i.e. the infection case fatality rate was 0.7% [8].

In 1996–2017, in Komi, in the European North of Russia, the TBE case fatality rate was 2.3% [28].

The incidence of both TBE and LD remains rather high in Northwestern Federal District of the Russian Federation (NWFDRF) [1, 3, 10, 12, 49]. For example, from 2006–2017 in Komi, the average TBE incidence rate ran to 1.4, while that of LD was 0.8 [1].

HGA, HME and Q fever are practically not recorded in the European north of Russia, although in some cases they are responsible for irreversible damage to the internal organs and disability of patients [5].

The choice of territories under study of human sera derives from the northward spread of ixodid ticks in the European north of Russia [10]. Such expansion in the tick's geographic range contributes to favorable conditions for TBDs distribution.

The objective of our work was to study the seroprevalence of IgG antibodies against TBEV, *Borrelia burgdorferi sensu lato*, *Coxiella burnetii*, *Anaplasma phagocytophilum*, and *Ehrlichia chaffeensis/E. muris* in the healthy population of NWFDRF: in the Arkhangelsk Oblast (AO), Leningrad Oblast (LO), Pskov Oblast (PO), Republic of Komi and Republic of Karelia. The purpose was to study the variation in seroprevalence of these indicators in different territories and whether there are age and gender differences.




Figure. Administrative divisions of the Northwestern Federal District

Note. 1 — Arkhangelsk Oblast, 2 — Vologda Oblast, 3 — Kaliningrad Oblast, 4 — Republic of Karelia, 5 — Komi Republic, 6 — Leningrad Oblast, 7 — Murmansk Oblast, 8 — Nenets Autonomous District, 9 — Novgorod Oblast, 10 — Pskov Oblast, 11 — St. Petersburg. The surveyed divisions (1, 4, 5, 6, 10) are highlighted in dark grey.

Table 1. Distribution of donors by gender, by age, and by territory of residence

Territory of residence	Number of donors	Gender		Age (years)					
		Male	Female	< 18	18–29	30–39	40–49	50–59	> 60
Republic of Komi	659	246	413	3	62	108	126	150	210
Republic of Karelia	292	148	144	0	35	109	95	45	8
AO	103	24	79	5	12	16	23	19	28
PO	98	42	56	0	21	23	35	17	2
LO	92	36	56	50	18	24	0	0	0
Total	1244	496	748	58	148	280	279	231	248

Materials and methods

The blood serum of healthy donors was sampled in 5 administrative divisions of NWFDRF: Komi, Karelia, AO, LO, PO. The territory of NWFDRF involves numerous islands, however, all samples were collected in mainland territory only. Figure shows a map of NWFDRF with the designation of its administrative units.

Blood was sampled in 2018–2019 after informed voluntary consent of donors and based on their personal data (Table 1). The inclusion criteria were as follows: good health at study entry; no professional risk of TBD acquisition; no history of vaccination against TBD and/or other infections caused by flaviviruses; no manifest forms of TBD; no diseases or conditions that affect the results of serological studies; residence in the territory under study and no trips out of it for several years at least.

Blood was sampled from the cubital vein into vacuum tubes, followed by centrifugation (10 min,

3000 rpm), freezing and storage of sera at -70°C . Subsequent transportation of the material was carried out in compliance with the “cold chain”. All sera samples were analyzed by ELISA with commercial test systems for detection of:

- IgG antibodies to TBEV — “VectoVKE-IgG” (CJSC “Vector-Best”, Novosibirsk),
- IgG antibodies to *Borrelia burgdorferi sensu lato* — “LimeBest” (CJSC “Vector-Best”, Novosibirsk),
- IgG antibodies to *Coxiella burnetii*, “Anti-Q” (Pasteur Institute, St. Petersburg),
- IgG antibodies to *Anaplasma phagocytophilum* — “GACH-ELISA-IgG” (LLC “Omnix”, St. Petersburg),
- IgG antibodies to *Ehrlichia chaffeensis*/*E. muris* — “MECH-IFA-IgG” (LLC “Omnix”, St. Petersburg).

All sera samples were tested according to the manufacturer’s manuals. The test results were classified as positive, negative, or borderline according

Table 2. IgG antibodies to TBD pathogens in sera of NWFDRF residents

Territory of residence	Number of samples	IgG-antibody-positive samples (number of positive/seroprevalence, %)						
		Antibody-positive to any TBD pathogens	Tick-borne encephalitis virus	<i>Borrelia burgdorferi sensu lato</i>	<i>C. burnetii</i>	<i>Anaplasma phagocytophilum</i>	<i>E. chaffeensis</i> / <i>E. muris</i>	Antibodies against 2 or more TBD pathogens
Republic of Komi	659	149/22.6	83/12.6	21/3.2	23/3.5	15/2.3	7/1.1**	15/2.3
Republic of Karelia	292	59/20.2	38/13.0	9/3.1	9/3.1	1/0.3	2/0.7	5/1.7
AO	103	15/14.6	7/6.8	2/1.9*	1/1.0	0	5/4.9**	1/1.0
PO	98	19/19.4	11/11.2	5/5.1	0	0	3/3.1	2/2.0
LO	92	28/30.4	13/14.1	6/6.5*	3/3.3	4/4.3	2/2.2	3/3.3
Total	1244	270/21.7	152/12.2	43/3.5	36/2.9	20/1.6	19/1.5	26/2.1

Note. When comparing the seroprevalence rates of IgG antibodies against TBD pathogens significant differences were found: *in relation to *Borrelia burgdorferi sensu lato* in AO and LO: the corresponding value of Fisher’s criterion is 1.651 (while for $p = 0.01$ the critical value is 1.450); **in relation to *E. chaffeensis*/*E. muris* in Komi and AO: the corresponding Fisher’s criterion is 2.244 (while the critical value for $p = 0.01$ is 1.450).

to the instructions enclosed in the diagnostic systems. Considering the probability of false positive results due to the common antigenic sites of some antigens, any borderline results were regarded as negative. In addition, for a more objective assessment of the results on TBEV, all sera containing IgG antibodies to TBEV according to VectoVKE-IgG system (CJSC Vector-Best, Novosibirsk, Russia) were simultaneously studied using the test system Enzygnost® Anti-TBE/FSME Virus (IgG) (Siemens Healthcare, GmbH, Marburg, Germany), both with specificity 99.5% according to the manufacturer's manual.

The distribution of our sera donors' by gender, age and territory of residence is shown in Table 1.

Results

In this study 1244 sera samples were tested for TBD pathogens. The parallel studies of 152 sera samples analysed for IgG antibodies to TBEV by VectoVKE-IgG were confirmed by Enzygnost® Anti-TBE Virus (IgG). The results of detection of IgG antibodies to the TBD pathogens *Borrelia burgdorferi sensu lato*, *Coxiella burnetii*, *Anaplasma phagocytophilum*, and *Ehrlichia chaffeensis/E. muris* are shown in Table 2.

From the 1244 studied sera samples IgG antibodies to TBD pathogens were detected in 270 (21.7%), where 26 (2.1%) samples contained antibodies to more than one pathogen. The overall seroprevalence was highest in LO (30.4%), then followed by Komi (22.6%), Karelia (20.2%), PO (19.4%) and AO (14.6%). IgG antibodies to all five TBD pathogens (TBEV, *Borrelia burgdorferi sensu lato*, *C. burnetii*, *Anaplasma phagocytophilum* and *E. chaffeensis/E. muris*) were found in sera samples from Komi, Karelia and LO. In AO we revealed antibodies to four (TBEV, *Borrelia burgdorferi sensu lato*, *C. burnetii* and *E. chaffeensis/E. muris*), while in PO only to three of them (TBEV, *Borrelia burgdorferi sensu lato* and *E. chaffeensis/E. muris*).

The number of samples from each territory was not sufficient to conclude on statistically significant differences in frequency of antibodies to the pathogens in this study.

However, from Table 2 the seroprevalence of IgG antibodies to TBEV and *Borrelia burgdorferi sensu lato* was 2.1 and 3.4 times higher in LO than in AO, respectively, while in Komi, Karelia and PO the percent of antibody-positive samples for these pathogens essentially did not differ, except for a slightly higher prevalence of *Borrelia* in PO and LO. In other territories the seroprevalence of IgG antibodies to TBEV did not vary significantly.

The highest seroprevalence to *A. phagocytophylum* was recorded in LO, to *E. chaffeensis/E. muris* the highest prevalence was detected in AO, and for *C. burnetii* IgG antibody seroprevalence in Karelia, Komi and LO was nearly the same.

IgG antibodies to all studied TBD pathogens were detected more often in men (55.2%) than in women (44.8%). The difference is especially demonstrated for IgG antibodies to TBEV where seroconversion was 15.7% in men, while only 9.9% in women.

In our study IgG antibodies to TBD did not depend considerably on the donor's age, being nearly the same for juniors (< 18 years) and seniors (> 60 years). However, IgG antibodies to *B. burgdorferi sensu lato* and *A. phagocytophylum* were detected in juniors somewhat more often than in seniors. IgG antibodies to *Coxiella burnetii* were highest in seniors (> 50 years), while IgG antibodies to *Borrelia* and *Anaplasma*, on the contrary, were highest in the youngest group (< 18 years) (see Table 3).

Discussion

Ixodid ticks are among the most important vectors of human TBDs, including TBE, LD, HGA and HME [4, 30]. The epidemiological profiles of those TBDs are nearly similar. Therefore, in the majority of cases IgG antibodies against TBEV, *B. burgdorferi sensu lato*, *A. phagocytophylum*, and *E. chaffeensis/E. muris* are indicative of frequent exposure to infected ticks within the studied territories. Seroprevalence is indicative of both the activity in TBD natural foci and the risk of infection in the local population [5]. On the contrary, the probability of catching Q fever directly from ixodid ticks is insignificant [25]; therefore, the IgG antibodies against

Table 3. Age distribution of donors with IgG antibodies to TBD pathogens

TBD pathogen	IgG antibody-positive/all donors; percent of antibody-positive (%)						
	< 18 n = 58	18–29 n = 148	30–39 n = 280	40–49 n = 279	50–59 n = 231	> 60 n = 248	Total n = 1244
TBEV	6/10.3%	19/12.8%	39/13.9%	39/14.0%	24/10.4%	25/10.1%	152/12.2%
<i>B. burgdorferi sensu lato</i>	4 /6.9%	3/2.0%	11/3.9%	8/2.9%	8/3.5%	9/3.6%	43/3.5%
<i>A. phagocytophilum</i>	3/5.2%	2/1.4%	1/0.4%	4/1.4%	4/1.7%	6 /2.4%	20/1.6%
<i>E. chaffeensis/E. muris</i>	2/3.4%	5/3.4%	1/0.4%	4/1.4%	3/1.3%	4/1.6%	19/1.5%
<i>C. burnetii</i>	1/1.7%	4/2.7%	5/1.8%	6/2.2%	9/3.9%	11/4.4%	36/2.9%

C. burnetii are more indicative of Q fever anthroponogenic foci activity. The role of ixodid ticks in the *C. burnetii* distribution is largely due to the infection of wild, agricultural and domestic animals. In addition, these blood-sucking arthropods infected with *C. burnetii* feed on migratory birds that transfer the infection to virgin territories [17, 52].

Here we present our first comprehensive serosurvey of humans in five administrative units of the NWFDRF for the presence of IgG antibodies against TBEV, *B. burgdorferi sensu lato*, *A. phagocytophilum* and *E. chaffeensis/E. muris* and *C. burnetii*. The results indicate that the proportion of TBD-infected populations in different territories varies.

Our results on the seroprevalence of IgG antibody to TBEV in AO (6.8%) and in Komi (12.6%) are close to those reported earlier (7.9% and 13.7%, respectively) [10, 51]. The index varies much depending on the territory under study. E.g., in the south of AO, it reached 20.9% [10]. Seroprevalence being the result of a permanent dynamic process depends both on the activity of TBD natural foci and the frequency of humans' exposure. An illustrative example is the increased seroprevalence of IgG antibodies to TBEV in the population of Komi between 2001 and 2013 from 3.5% to 13.7%, probably due to climatic changes [49], that favor both northward expanding of tick geographic ranges and exposure of local inhabitants to infected tick bites [51]. An increase in TBEV seroprevalence is typical not only for the North of Russia; a similar trend has been observed in some regions of the Czech Republic [35].

Here we studied the seroprevalence of IgG antibodies to TBEV in the healthy population of the Northwestern Russia, while researchers in Europe focus mostly on occupationally-risk groups. However, in many European countries the reported TBEV-seroprevalence in high risk occupationally exposed groups is much less (e.g., only 2.3% in France [43]), than that of healthy donors in NWFDRF, where the seroprevalence is 12.2%. Serosurveillance of 1213 healthy donors in Western Norway showed little to no TBEV prevalence among local residents [31]. Similar results were obtained from a survey of healthy donors in other areas (Vestfold and Telemark counties) in Norway. Only 0.4% of them had antibodies to TBEV [36, 38].

Our findings on seroprevalence of IgG antibodies to *B. burgdorferi sensu lato*. In NWFDRF (3.5%) do not differ much from those reported in other European countries. E.g., in Belgium *B. burgdorferi sensu lato* seroprevalence in a healthy population of blood donors was 2.6–2.9% [21].

HGA must be classified as an emerging infection in NWFDRF. HGA was first diagnosed in 1990 in Wisconsin (USA) in a patient with a tick bite history and severe febrile illness [18]. However, under-diagnosis of HGA is typical for many coun-

tries, including USA [40]. The incidence of human HGA cases in Europe is lower (estimated under 300) than reported from the USA, where a steady increase has been reported since 2001, with more than 15,000 accumulated cases until 2015 [16]. HGA cases have now been reported from Austria, Croatia, France, Italy, Latvia, the Netherlands, Norway, Poland, Slovenia, Spain, and Sweden [37, 39]. There is an incongruence between human seroprevalence and observed clinical cases that might arise from incomplete diagnosis, or a high rate of asymptomatic infections [41]. In Russia, as well as in other European countries, HGA is rarely registered. In many territories of Russia, where ticks infected with *A. phagocytophilum* were found, HGA is not diagnosed [9]. A similar situation is observed in NWFDRF, where *A. phagocytophilum* was also found in ticks [26], and HGA is not diagnosed in humans either. It is possible that HGA patients are given a different diagnosis, for example, "Flu" or "fever of unknown etiology." The discrepancy between the relatively high seroprevalence rates for *A. phagocytophilum* (4.3% in LO), and the lack of diagnosis of this infection in humans in NWFDRF, revealed by us, is probably due to the wide spread of asymptomatic forms of this disease and the imperfection of its diagnosis [41]. Despite the obviously ubiquitous presence of *A. phagocytophilum* in ticks and various wild and domestic animals in the European North of Russia [26], the reported HGA clinical cases are very rare, and territories with *A. phagocytophilum*-infected ticks are much more numerous than those with a few registered HGA cases [41]. However, in S.P. Botkin hospital in St. Petersburg antibodies to HGA were detected in 16 of 934 tick-bitten febrile patients, while no other antibodies to TBD pathogens (TBEV or *B. burgdorferi sensu lato*) were found. In two of these 16 patients an increase in IgG antibodies to *A. phagocytophilum* was recorded which, along with the clinical picture and pre-existing tick bites, made it possible to diagnose HGA. In Europe rather high HGA seroprevalence rate was detected in TBE-suspected patients: 30.96% in Belgium [20], 25% in Slovakia [34], 16.28% in donors in Norway [31], 11.4–28% in Sweden [40, 46], and 3.5–9.4% in Poland [19, 53]. According to our findings the average level of HGA seroprevalence in the healthy population of NWFDRF was 1.6% (though 4.3% in LO) that is less than in most of the European countries mentioned above. This comparatively low seroprevalence, on the one hand, may stem from lower *A. phagocytophilum* prevalence in the territories under study; on the other hand, it may be due to different methods of antibodies detection and differences in the surveyed contingents. To detect IgG antibodies to *A. phagocytophilum* in Europe the IFA assay was mostly applied to survey "occupational

risk” groups, while here we used ELISA and applied it in healthy donors without “occupational risk”. However, in our opinion, even the relatively low figures for *A. phagocytophilum* seroprevalence in LO, Komi and Karelia substantiate the need to examine tick-bitten febrile patients for the presence of antibodies to *A. phagocytophilum*.

HME, despite its clinical resemblance to HGA, is characterized by more severe course and a higher mortality rate [24]. HME is widespread in the southeastern, southern, central, and mid-Atlantic parts of the United States. In 2015, *E. chaffeensis* infection was reported in 35 states, with most numerous cases in Missouri, Arkansas, New York, and Virginia. In the European Union, *E. chaffeensis* is not subject to registration [37], however, HME cases have been reported in Serbia [15], Czech Republic, Denmark, Sweden, and other countries [28], but those findings are fragmentary. In the Russian regions diagnosis of HME cases are rare compared to detection of *E. chaffeensis/E. muris* in ticks [1]. However, our earlier serological studies revealed the HME occurrence in the North-West of Russia: e.g., in the healthy population of St. Petersburgs outskirts the seroprevalence of IgG antibodies to *E. chaffeensis/E. muris* was 4.4% [50].

In this study the seroprevalence of IgG antibodies to *E. chaffeensis/E. muris* in average was 1.5%. It was revealed in all territories under study and varied from 0.7% (in Karelia) to 4.9% (AO). Those findings are consistent with an earlier communication on detection of IgG antibodies to *E. chaffeensis/E. muris* in the sera of 23.3% of febrile patients hospitalized in Arkhangelsk city after tick bites [50].

Q fever is diagnosed in humans and animals, and its natural or household foci are identified practically all over the world, with the possible exception for New Zealand [14]. Not all countries have mandatory registration of Q fever incidence, however, the infection is usually diagnosed where appropriate research is carried out [33]. In Europe, Q fever is regularly notified in many countries, e.g., in Bulgaria, France and Germany [27]. The largest known Q fever outbreak reported to date involved approximately 4,000 human cases (25 fatal) and occurred during 2007–2010 in the Netherlands [22, 25, 44]. In the NWFDRF, a large outbreak of Q fever was recorded at the end of 20th century [2].

In our study, IgG antibodies to *C. burnetii* were detected in all studied territories, with the exception for PO, while in Karelia, Komi and LO it was rather high (> 3%).

However, Q fever is not diagnosed in NWFDRF in recent years, and only one Q fever case was diagnosed in LO in 2017–2019.

Among the reasons for Q fever under-diagnosis is the difficulty of its clinical recognition due to pronounced polymorphism of the disease manifestations and the absence of pathognomonic symptoms.

Significant under-diagnosis of Q fever causes irrational treatment, resulting in chronicity. Chronic Q fever is a very frequent outcome of the disease. For example, in the south of France 5–8% of all endocarditis cases are caused by *C. burnetii* [48]. Such development of the disease in humans is often fatal [42].

Comparison of IgG antibody seroprevalence in LO (3.3%) with the same long-term indicator in the city of St. Petersburg (0.4%) [6] lends indirect support to the view that ixodid ticks play a relatively small role in *C. burnetii* infection in humans. The overwhelming majority of St. Petersburg residents annually spend their recreation time in LO and tick-bitten by ixodid ticks [12]. However, the IgG antibody seroprevalence in the city residents is much lower than in the LO population, because a larger part of the LO population tend to temporary be in contact with animals at livestock enterprises or private farms compared to city residents.

In our study, IgG antibodies to TBDs were detected in humans of various ages, from children (< 18 years) to the elderly (> 60 years). Similar findings are available for Russia as a whole [7]. The age dependence of seroprevalence was insignificant.

The most significant gender differences in seroprevalence rates were related to TBE.

In men, the seroprevalence of IgG antibodies to TBEV exceeded that in women, probably to a certain extent due to their exposure to ticks during traditional out-of-doors occupations (fishing, hunting, hiking, etc.).

Analysis of modern literature on infections transmitted by ixodid ticks suggests that any disease resulting from tick bites should be considered as a potential mixed infection. Coinfection of two or more pathogens during the disease development complicates the diagnosis of mixed infections [4]. It was found that in northwestern Russia, mixed infection without pronounced clinical manifestations occurs 7.6 times more often than with the development of the clinical picture of diseases [11, 13]. In our study, antibodies to two or more pathogens were detected in 2.1% of sera samples, and as a rule, those were IgG antibodies to TBEV and *Borrelia burgdorferi sensu lato*. Our findings do not allow us to determine whether people were infected with those two pathogens simultaneously or from different tick bites. Although, considering the pathogens ability to persist in a human body, it can be assumed that individuals with coinfections may have a greater risk of chronicity.

When testing sera by the ELISA methods one cannot completely exclude the possibility of false-positive results. To minimize this risk in our study we interpreted the borderline results as negative, and similar results from two different test systems in parallel was required to conclude that a sample was positive.

However, for our purpose, the absolute figures are less important than the variation in seroprevalence of those indicators in different territories, and in persons of different ages and gender.

Conclusions

A sizable portion (on average 21.7%) of NWFDRF population is infected with TBD pathogens. The percent of infected population varies significantly depending on the territory. The presence of IgG antibodies to TBD pathogens in the sera of humans in the absence of reported incidence is indicative of underdiagnosis of those TBDs. The discrepancy between our findings on seroprevalence and the reported TBDs prevalence is largely due to the absence of pathognomonic symptoms in these diseases, its mild, asymptomatic clinical course in some cases, and insufficient laboratory diagnostics. Significant underdiagnosis of TBDs as a consequence of irrational treatment results in chronic course and lethality, therapeutic misconception, erroneous ideas about the real distribution of TBDs and the lack of proper alertness of medical staff to those diseases.

References

- Гнатив Б.Р., Токаревич Н.К. Результаты многолетнего мониторинга клещевого вирусного энцефалита и клещевого боррелиоза в Республике Коми // Инфекция и иммунитет. 2021. Т. 11, № 4. С. 707–722. [Gnativ B.R., Tokarevich N.K. Long-term monitoring of tick-borne viral encephalitis and tick-borne borreliosis in the Komi Republic. *Infektsiya i imunitet = Russian Journal of Infection and Immunity*, 2021, vol. 11, no. 4, pp. 707–722. (In Russ.)] doi: 10.15789/2220-7619-ROL-1299
- Дайтер А.Б., Рыбакова Н.А., Токаревич Н.К., Самитова В.И., Лимин Б.В. Эпидемиологическая проекция внутристадных очагов лихорадки Ку // Журнал микробиологии, эпидемиологии и иммунобиологии. 1988. Т. 11. С. 51–56. [Daiter A.B., Rybakova N.A., Tokarevich N.K., Samitova V.I., Limin B.V. An epidemiologic projection of foci of Q fever within herds. *Zhurnal mikrobiologii, epidemiologii i immunobiologii = Journal of Microbiology, Epidemiology and Immunobiology*, 1988, vol. 11, pp. 51–56. (In Russ.)]
- Карташов М.Ю., Микрюкова Т.П., Кривошеина Е.И., Кузнецова А.И., Глушкина Л.И., Корабельников И.В., Егорова Ю.И., Терновой В.А., Локтев В.Б. Генотипирование возбудителей клещевого энцефалита и лихорадки Кемерово в таежных клещах, собранных в Республике Коми // Инфекция и иммунитет. 2020. Т. 10, № 1. С. 159–166. [Kartashov M.Yu., Mikryukova T.P., Krivosheina E.I., Kuznetsova A.I., Glushkova L.I., Korabel'nikov I.V., Egorova Yu.I., Ternovoi V.A., Loktev V.B. Genotyping of tick-borne encephalitis and Kemerovo viruses in taiga ticks collected in the Komi Republic. *Infektsiya i imunitet = Russian Journal of Infection and Immunity*, 2020, vol. 10, no. 1, pp. 159–166. (In Russ.)] doi: 10.15789/2220-7619-GOT-1147
- Конькова-Рейдман А.Б., Злобин В.И. Клинический полиморфизм иксодовых клещевых боррелиозов (микст-инфекция с клещевым энцефалитом) на территории Южно-Уральского региона России // Сибирский медицинский журнал. 2011. № 1. С. 17–19. [Kon'kova-Reidman A.B., Zlobin V.I. Clinical polymorphism of Ixodes tick-borne borrelioses (mixed infection with tick-borne encephalitis) on the territory of South-Ural Region of Russia. *Sibirskij medicinskij zurnal (Irkutsk) = Siberian Medical Journal (Irkutsk)*, 2011, vol. 1, pp. 17–19. (In Russ.)]
- Коренберг Э.И., Помелова В.Г., Осин Н.С. Природноочаговые инфекции, передающиеся иксодовыми клещами / Под ред. А.Л. Гинцбурга и В.Н. Злобина. М.: Наука, 2013. 464 с. [Korenberg E.I., Pomelova V.G., Osin N.S. Infections with natural focality transmitted by ixodid ticks. Eds: A.L. Ginzburg, V.N. Zlobin. Moscow: Nauka, 2013. 463 p. (In Russ.)]
- Краева Л.А., Токаревич Н.К., Лаврентьева И.Н., Рошина Н.Г., Кафтырева Л.А., Кунилова Е.С., Курова Н.Н., Стоянова Н.А., Антикова А.Ю., Сварваль А.В., Зуева Е.В., Порин А.А., Рогачева Е.В., Желтакова И.Р., Хамитова И.В., Тимофеева Е.В., Беспалова Г.И. Инфицированность трудовых мигрантов из Средней Азии и постоянных жителей Санкт-Петербурга возбудителями различных инфекционных заболеваний и восприимчивость к ним // Инфекция и иммунитет. 2018. Т. 8, № 1. С. 61–70. [Kraeva L.A., Tokarevich N.K., Lavrentyeva I.N., Roshchina N.G., Kaftyreva L.A., Kunilova E.S., Kurova N.N., Stoyanova N.A., Antikova A.Yu., Svarval A.V., Zueva E.V., Porin A.A., Rogacheva E.V., Zheltakova I.R., Khamitova I.V., Timofeeva E.V., Bespalova G.I. Infection of labour migrants from Central Asia and residents of St. Petersburg and their susceptibility to various infectious diseases. *Infektsiya i imunitet = Russian Journal of Infection and Immunity*, 2018, vol. 8, no. 1, pp. 61–70. (In Russ.)] doi: 10.15789/2220-7619-2018-1-61-70
- Никитин А.Я., Андаев Е.И., Носков А.К., Пакскина Н.Д., Яцменко Е.В., Веригина Е.В., Балахонов С.В. Особенности эпидемиологической ситуации по клещевому вирусному энцефалиту в Российской Федерации в 2017 г. и прогноз ее развития на 2018 г. // Проблемы особо опасных инфекций. 2018. Т. 1. С. 44–49. [Nikitin A.Ya., Andaev E.I., Noskov A.K., Pakskina N.D., Yatsmenko E.V., Verigina E.V., Balakhonov S.V. Epidemiological situation of tick-borne viral encephalitis in the Russian Federation in 2017 and forecast of its development in 2018. *Problemy osobopopasnykh infektsii = Problems of Special Pathogenic Diseases*, 2018, vol. 1, pp. 44–49. (In Russ.)]

Men are infected with TBDs more often than women. However, people, starting from children (< 18 years old) to the elderly (> 60 years old) were found to be infected with TBD pathogens almost evenly irrespective of age. Infection with two or more TBD pathogens was detected in 2.1% of the surveyed humans.

The detection of IgG antibodies to TBD pathogens in the northern territories, where TBDs never had been diagnosed, confirms indirectly the northward expansion of tick habitats [10, 46, 51]. Similar expansion of ixodid tick habitats has also been observed in other countries of the Northern Europe [32, 45, 47]. Of particular concern is the health of the indigenous population of the North, who historically had never been exposed to ticks and more vulnerable to tick-borne infections [29].

Acknowledgments

The project has been partially funded by the Norwegian Ministry of Health and Care services, Barents Program, Project number B1710.

- Pakskina N.D., Yatsmenko E.V., Verigina E.V., Balakhonov S.V. Peculiarities of the epidemiological situation on tick-borne viral encephalitis in the Russian Federation in 2017 and the forecast for 2018. *Problemy osobo opasnykh infektsii = Problems of Particularly Dangerous Infections*, 2018, vol. 1, pp. 44–49. (In Russ.) doi: 10.21055/0370-1069-2018-1-44-49
8. Носков А.К., Андаев Е.И., Никитин А.Я., Пакскина Н.Д., Яцменко Е.В., Веригина Е.В., Толмачева М.И., Балахонов С.В. Заболеваемость клещевым вирусным энцефалитом в субъектах Российской Федерации. Сообщение 1: Эпидемиологическая ситуация по клещевому вирусному энцефалиту в 2018 г. и прогноз на 2019 г. // Проблемы особы опасных инфекций. 2019. Т. 1. С. 74–80. [Noskov A.K., Andaev E.I., Nikitin A.Ya., Pakskina N.D., Yatsmenko E.V., Verigina E.V., Tolmacheva M.I., Balakhonov S.V. Tick-borne viral encephalitis morbidity rates in the constituent entities of the Russian Federation. Communication 1. Epidemiological situation on tick-borne viral encephalitis in 2018 and forecast for 2019. *Problemy osobo opasnykh infektsii = Problems of Particularly Dangerous Infections*, 2019, vol. 1, pp. 74–80. (In Russ.) doi: 10.21055/0370-1069-2019-1-74-80]
 9. Проворова В.В., Краснова Е.И., Хохлова Н.И., Савельева М.А., Филимонова Е.С., Кузнецова В.Г. Старые и новые клещевые инфекции в России // Инфекционные болезни: новости, мнения, обучение. 2019. Т. 8, № 2. С. 102–112. [Provorova V.V., Krasnova E.I., Khokhlova N.I., Savel'eva M.A., Filimonova E.S., Kuznetsova V.G. Tissue infections in Russia. *Infectsionnye bolezni: novosti, mneniya, obuchenie = Infectious Diseases: News, Opinions, Training*, 2019, vol. 8, no. 2, pp. 102–112. (In Russ.) doi: 10.24411/2305-3496-2019-12013]
 10. Соколова О.В., Чашин В.П., Попова О.Н., Бузинов Р.В., Пасынкова М.М., Гудков А.Б. Эпидемиологические особенности распространения клещевого вирусного энцефалита в Архангельской области // Экология человека. 2017. Т. 24, № 4. С. 12–19. [Sokolova O.V., Chashchin V.P., Popova O.N., Buzinov R.V., Pasynkova M.M., Gudkov A.B. Epidemiological character of tick-borne viral encephalitis extension in the Arkhangelsk Region. *Ekologiya cheloveka = Human Ecology*, 2017, vol. 4, pp. 12–19. (In Russ.) doi: 10.33396/1728-0869-2017-4-12-19]
 11. Субботина Н.С., Доршакова Н.В., Петрова А.В. Эпидемиологическая характеристика клещевого энцефалита в Северо-Западном регионе России // Экология человека. 2007. № 7. С. 15–19. [Subbotina N.S., Dorshakova N.S., Petrova A.V. Epidemiological characteristic of tick-borne encephalitis in North-West region of Russia. *Ekologiya cheloveka = Human Ecology*, 2007, vol. 7, pp. 15–19. (In Russ.)]
 12. Сюзюмова Е.А., Тельнова Н.В., Шапарь А.О., Асланов Б.И., Стоянова Н.А., Токаревич Н.К. Эколо-эпидемиологическая характеристика клещевого энцефалита в Санкт-Петербурге // Инфекция и иммунитет. 2020. Т. 10, № 3. С. 533–542. [Suziumova E.A., Telnova N.V., Shapar A.O., Aslanov B.I., Stoyanova N.A., Tokarevich N.K. Ecological and epidemiological characteristics of tick-borne encephalitis in St. Petersburg. *Infektsiya i immunitet = Russian Journal of Infection and Immunity*, 2020, vol. 10, no. 3, pp. 533–542. (In Russ.) doi: 10.15789/2220-7619-EAE-924]
 13. Усков А.Н., Байгеленов К.Д., Бургасова О.А., Гринченко Н.Е. Современные представления о диагностике клещевых инфекций // Сибирский медицинский журнал (Иркутск). 2008. Т. 7. С. 148–152. [Uskov A.N., Baygelenov K.D., Burgasova O.A., Grintchenko N.E. Present view on diagnostics of tick-borne infections. *Sibirskij medicinskij zurnal (Irkutsk) = Siberian Medical Journal (Irkutsk)*, 2008, vol. 7, pp. 148–152. (In Russ.)]
 14. Angelakis E., Raoult D. Q Fever. *Vet. Microbiol.*, 2010, vol. 140, no. 3–4, pp. 297–309. doi: 10.1016/j.vetmic.2009.07.016
 15. Arsić B., Gligić A., Ristanović E., Lako B., Potkonjak A., Perunicić M., Pavlović M. A case of human monocytic ehrlichiosis in Serbia. *Srp. Arh. Celok. Lek.*, 2014, vol. 142, no. 1–2, pp. 79–82. doi: 10.2298/sarh1402079a
 16. Bakken J.S., Dumler J.S. Human granulocytic anaplasmosis. *Infect. Dis. Clin. North Am.*, 2015, vol. 29, no. 2, pp. 341–355. doi: 10.1016/j.idc.2015.02.007
 17. Berthová L., Slobodník V., Slobodník R., Olekšák M., Sekeyová Z., Svitálková Z., Kazimírová M., Špitalská E. The natural infection of birds and ticks feeding on birds with *Rickettsia* spp. and *Coxiella burnetii* in Slovakia. *Exp. Appl. Acarol.*, 2016, vol. 68, no. 3, pp. 299–314. doi: 10.1007/s10493-015-9975-3
 18. Chen S.M., Dumler J.S., Bakken J.S., Walker D.H. Identification of a granulocytotropic *Ehrlichia* species as the etiologic agent of human disease. *J. Clin. Microbiol.*, 1994, vol. 32, no. 3, pp. 589–595. doi: 10.1128/jcm.32.3.589-595.1994
 19. Chmielewska-Badora J., Moniuszko A., Żukiewicz-Sobczak W., Zwoliński J., Piątek J., Pancewicz S. Serological survey in persons occupationally exposed to tick-borne pathogens in cases of co-infections with *Borrelia burgdorferi*, *Anaplasma phagocytophilum*, *Bartonella* spp. and *Babesia microti*. *Ann. Agric. Environ. Med.*, 2012, vol. 19, no. 2, pp. 271–274.
 20. Cochez C., Ducofre G., Vandenvelde C., Luyasu V., Heyman P. Human anaplasmosis in Belgium: a 10-year seroepidemiological study. *Ticks Tick Borne Dis.*, 2011, vol. 2, pp. 156–159. doi: 10.1016/j.ttbdis.2011.06.004
 21. De Keukeleire M., Vanwambeke S.O., Cochez C., Heyman P., Fretin D., Deneys V., Luyasu V., Kabamba B., Robert A. Seroprevalence of *Borrelia burgdorferi*, *Anaplasma phagocytophilum*, and *Francisella tularensis* infections in Belgium: results of three population-based samples. *Vector Borne Zoonotic. Dis.*, 2017, vol. 17, no. 2, pp. 108–115. doi: 10.1089/vbz.2016.1954
 22. De Rooij M.M., Borlée F., Smit L.A., de Bruin A., Janse I., Heederik D.J., Wouters I.M. Detection of *Coxiella burnetii* in ambient air after a large Q fever outbreak. *PLoS One*, 2016, vol. 11, no. 3: e0151281. doi: 10.1371/journal.pone.0151281
 23. Dumler J.S., Dotevall L., Gustafson R., Granström M. A population-based seroepidemiologic study of human granulocytic ehrlichiosis and Lyme borreliosis on the west coast of Sweden. *J. Infect. Dis.*, 1997, vol. 175, no. 3, pp. 720–722. doi: 10.1093/infdis/175.3.720
 24. Dumler J.S., Madigan J.E., Pusterla N., Bakken J.S. Ehrlichioses in humans: epidemiology, clinical presentation, diagnosis, and treatment. *Clin. Infect. Dis.*, 2007, vol. 45, suppl. 1: S45–51. doi: 10.1086/518146
 25. Eldin C., Mélenotte C., Mediannikov O., Ghigo E., Million M., Edouard S., Mege J.L., Maurin M., Raoult D. From Q fever to *Coxiella burnetii* infection: a paradigm change. *Clin. Microbiol. Rev.*, 2017, vol. 30, no. 1, pp. 115–190. doi: 10.1128/CMR.00045-16
 26. Eremeeva M.E., Oliveira A., Moriarity J., Robinson J.B., Tokarevich N.K., Antyukova L.P., Pyanyh V.A., Emeljanova O.N., Ignatjeva V.N., Buzinov R., Pyankova V., Dasch G.A. Detection and identification of bacterial agents in *Ixodes persulcatus*

- Schulze ticks from the north western region of Russia. *Vector Borne Zoonotic Dis.*, 2007, vol. 7, no. 3, pp. 426–436. doi: 10.1089/vbz.2007.0112
27. Georgiev M., Afonso A., Neubauer H., Needham H., Thiery R., Rodolakis A., Roest H., Stark K., Stegeman J., Vellema P., van der Hoek W., More S. Q fever in humans and farm animals in four European countries, 1982 to 2010. *Euro Surveill.*, 2013, vol. 18, no. 8: 20407.
 28. Gratz N.G., World Health Organization. The vector-borne human infections of Europe: their distribution and burden on public health. *Copenhagen: WHO Regional Office for Europe*, 2004. 144 p.
 29. Hedlund C., Blomstedt Y., Schumann B. Association of climatic factors with infectious diseases in the Arctic and subarctic region — a systematic review. *Glob. Health Action*, 2014, vol. 7: 24161. doi: 10.3402/gha.v7.24161
 30. Heyman P., Cochez C., Hofhuis A., van der Giessen J., Sprong H., Porter S.R., Losson B., Saegerman C., Donoso-Mantke O., Niedrig M., Papa A. A clear and present danger: tick-borne diseases in Europe. *Expert Rev. Anti Infect. Ther.*, 2010, vol. 8, no. 1, pp. 33–50. doi: 10.1586/eri.09.118
 31. Hjetland R., Henningsson A.J., Vainio K., Dudman S.G., Grude N., Ulvestad E. Seroprevalence of antibodies to tick-borne encephalitis virus and Anaplasma phagocytophilum in healthy adults from western Norway. *Infect. Dis. (Lond.)*, 2015, vol. 47, no. 1, pp. 52–56. doi: 10.3109/00365548.2014.959044
 32. Jaenson T.G., Hjertqvist M., Bergstrom T., Lundkvist A. Why is tick-borne encephalitis increasing? A review of the key factors causing the increasing incidence of human TBE in Sweden. *Parasit Vectors*, 2012, vol. 5: 184. doi: 10.1186/1756-3305-5-184
 33. Kazar J. Coxiella burnetii infection. *Ann. NY Acad. Sci.*, 2005, vol. 1063, pp. 105–114. doi: 10.1196/annals.1355.018
 34. Kocianová E., Košt'ánová Z., Štefanidesová K., Špitalská E., Boldiš V., Hučková D., Stanek G. Serologic evidence of Anaplasma phagocytophilum infections in patients with a history of tick bite in central Slovakia. *Wien Klin. Wochenschr.*, 2008, vol. 120, pp. 427–431. doi: 10.1007/s00508-008-1000-y
 35. Kříž B., Kott I., Daniel M., Vráblík T., Beneš Č. Impact of climate changes on the incidence of tick-borne encephalitis in the Czech Republic in 1982–2011. [In *Czech.J. Epidemiol. Mikrobiol. Imunol.*, 2015, vol. 64, pp. 24–32.]
 36. Larsen A.L., Kanestrøm A., Bjørland M., Andreassen A., Soleng A., Vene S., Dudman S.G. Detection of specific IgG antibodies in blood donors and tick-borne encephalitis virus in ticks within a non-endemic area in southeast Norway. *Scand. J. Infect. Dis.*, 2014, vol. 46, no. 3, pp. 181–184. doi: 10.3109/00365548.2013.865140
 37. Madison-Antenucci S., Kramer L.D., Gebhardt L.L., Kauffman E. Emerging tick-borne diseases. *Clin. Microbiol. Rev.*, 2020, vol. 33, no. 2: e00083-18. doi: 10.1128/CMR.00083-18
 38. Marvik A., Tveten Y., Pedersen A.-B., Stiasny K., Andreassen A.K., Grude N. Low prevalence of tick-borne encephalitis virus antibodies in Norwegian blood donors. *Infect. Dis.*, 2021, vol. 53, no. 1, pp. 44–51. doi: 10.1080/23744235.2020.1819561
 39. Matei I.A., Estrada-Peña A., Cutler S.J., Vayssier-Taussat M., Varela-Castro L., Potkonjak A., Zeller H., Mihalca A.D. A review on the eco-epidemiology and clinical management of human granulocytic anaplasmosis and its agent in Europe. *Parasit. Vectors*, 2019, vol. 12: 599. doi: 10.1186/s13071-019-3852-6
 40. Mendell N.L., Reynolds E.S., Blanton L.S., Hermance M.E., Londoño A.F., Hart C.E., Quade B.R., Esterly A.T., Hendrix C.B., Teel P.D., Bouyer D.H., Thangamani S. Detection of Rickettsiae, Borreliae, and Ehrlichiae in ticks collected from Walker County, Texas, 2017–2018. *Insects*, 2019, vol. 10, no. 10: 315. doi: 10.3390/insects10100315
 41. Nordberg M. Tick-borne infections in humans: aspects of immunopathogenesis, diagnosis and co-infections with *Borrelia burgdorferi* and *Anaplasma phagocytophilum*. *Sweden: Linköping University*, 2012.
 42. Raoult D., Tissot-Dupont H., Foucault C., Gouvernet J., Fournier P.E., Bernit E., Stein A., Nesri M., Harle J.R., Weiller P.J. Q fever 1985–1998. Clinical and epidemiologic features of 1,383 infections. *Medicine*, 2000, vol. 79, no. 2, pp. 109–123. doi: 10.1097/00005792-200003000-00005
 43. Rigaud E., Jaulhac B., Garcia-Bonnet N., Hunfeld K.P., Féminé F., Huet D., Goulvestre C., Vaillant V., Deffontaines G., Abadia-Benoist G. Seroprevalence of seven pathogens transmitted by the *Ixodes ricinus* tick in forestry workers in France. *Clin. Microbiol. Infect.*, 2016, vol. 22, no. 8: 735.e1-9. doi: 10.1016/j.cmi.2016.05.014
 44. Schimmer B., Dijkstra F., Vellema P., Schneeberger P.M., Hackert V., ter Schegget R., Wijkmans C., van Duynhoven Y., van der Hoek W. Sustained intensive transmission of Q fever in the south of the Netherlands, 2009. *Euro Surveill.*, 2009, vol. 14, no. 19: 19210. doi: 10.2807/ese.14.19.19210-en
 45. Skarpaas T., Golovljova I., Vene S., Ljøstad U., Sjursen H., Plyusnin A., Lundkvist A. Tick borne encephalitis virus, Norway and Denmark. *Emerg. Infect. Dis.*, 2006, vol. 12, pp. 1136–1138. doi: 10.3201/eid1207.051567
 46. Skarpaas T., Sundøy A., Bruu A.L., Vene S., Pedersen J., Eng P.G., Csányi P.A. Skogflåttencefalitt i Norge [Tick-borne encephalitis in Norway]. *Tidsskrift for den Norske laegeforening: tidsskrift for praktisk medicin, ny række*, 2002, vol. 122, no. 1, pp. 30–32.
 47. Süss J. TBE — a short overview on epidemiological status in Europe. *ISW-TBE: Vienna, Austria*, 2012; Feb. 2–3.
 48. Tissot Dupont H., Raoult D., Brouqui P., Janbon F., Peyramond D., Weiller P.J., Chicheportiche C., Nezri M., Poirier R. Epidemiologic features and clinical presentation of acute Q fever in hospitalized patients: 323 French cases. *Am. J. Med.*, 1992, vol. 93, no. 4, pp. 427–434. doi: 10.1016/0002-9343(92)90173-9
 49. Tokarevich N., Stoyanova N., Gnativ B., Kazakovtsev S., Blinova O., Revich B. Seroprevalence of tick-borne diseases in the population of the European North of Russia. *Med. Safety Glob. Health*, 2017, vol. 6: 132. doi: 10.4172/2574-0407.1000132
 50. Tokarevich N., Stoyanova N., Gracheva L., Shulaykina I., Kulikov A., Kozarenko A. Lyme Borreliosis and other Tick-borne diseases. *10th Int. Conference. Vienna, Austria*, 2005, p. 48.
 51. Tokarevich N., Tronin A., Gnativ B., Revich B., Blinova O., Evengard B. Impact of air temperature variation on the ixodid ticks habitat and tick-borne encephalitis incidence in the Russian Arctic: the case of the Komi Republic. *Int. J. Circumpolar Health*, 2017, vol. 76, no. 1: 1298882. doi: 10.1080/22423982.2017.1298882
 52. Tokarevich N.K., Panferova Y.A., Freylikhman O.A., Blinova O.V., Medvedev S.G., Mironov S.V., Grigoryeva L.A., Tretyakov K.A., Dimova T., Zaharieva M.M., Nikolov B., Zehtindjiev P., Najdenski H. Coxiella burnetii in ticks and wild birds. *Ticks & Tick Borne Dis.*, 2019, vol. 10, no. 2, pp. 377–385. doi: 10.1016/j.ttbdis.2018.11.020

53. Walory J., Bukowska B., Grzesiowski P., Czarnecka I., Paluchowska E., Zabielski S., Grzywocz A. Wystepowanie przeciwiał przeciw Anaplasma phagocytophilum, Babesia microti i Borrelia burgdorferi u dorosłych z północno-wschodnich rejonów Polski [Prevalence of antibodies against Anaplasma phagocytophilum, Babesia microti i Borrelia burgdorferi in adults in North-Eastern Poland]. *Polski merkuriusz lekarski: organ Polskiego Towarzystwa Lekarskiego*, 2005, vol. 19, no. 114, pp. 754–757.
54. Wittesjö B., Björsdorff A., Eliasson I., Berglund J. First long-term study of the seroresponse to the agent of human granulocytic ehrlichiosis among residents of a tick-endemic area of Sweden. *Eur. J. Clin. Microbiol. Infect. Dis.*, 2001, vol. 20, no. 3, pp. 173–178. doi: 10.1007/s100960100463

Авторы:

Токаревич Н.К., д.м.н., профессор, зав. лабораторией зоононтропонозных инфекций ФБУН НИИ эпидемиологии и микробиологии имени Пастера, Санкт-Петербург, Россия;
Блинова О.В., к.х.н., младший научный сотрудник лаборатории зоононтропонозных инфекций ФБУН НИИ эпидемиологии и микробиологии имени Пастера, Санкт-Петербург, Россия;
Стоянова Н.А., к.м.н., ведущий научный сотрудник лаборатории зоононтропонозных инфекций ФБУН НИИ эпидемиологии и микробиологии имени Пастера, Санкт-Петербург, Россия;
Баймова Р.Р., младший научный сотрудник лаборатории зоононтропонозных инфекций ФБУН НИИ эпидемиологии и микробиологии имени Пастера, Санкт-Петербург, Россия;
Сюзюмова Е.А., младший научный сотрудник лаборатории зоононтропонозных инфекций ФБУН НИИ эпидемиологии и микробиологии имени Пастера, Санкт-Петербург, Россия;
Ломоносова В.И., младший научный сотрудник лаборатории зоононтропонозных инфекций ФБУН НИИ эпидемиологии и микробиологии имени Пастера, Санкт-Петербург, Россия;
Тронин А.А., д.г.-м.н., директор Санкт-Петербургского научно-исследовательского центра экологической безопасности РАН, Санкт-Петербург, Россия;
Бузинов Р.В., д.м.н., доцент, зам. директора по развитию Федеральной службы по надзору в сфере защиты прав потребителей и благополучия человека по Архангельской области, г. Архангельск, Россия;
Соколова О.В., зам. начальника отдела эпидемиологического надзора Управления Роспотребнадзора по Архангельской области, г. Архангельск, Россия;
Гнатив Б.Р., зам. начальника отдела эпидемиологического надзора Управления Роспотребнадзора по Республике Коми, г. Сыктывкар, Россия;
Буц Л.В., начальник отдела эпидемиологического надзора Управления Роспотребнадзора по Ленинградской области, Санкт-Петербург, Россия;
Бубнова Л.А., зав. эпидемиологическим отделом ФБУЗ Центр гигиены и эпидемиологии в Республике Карелия, г. Петрозаводск, Россия;
Сафонова О.С., биолог ФБУЗ Центр гигиены и эпидемиологии в Республике Карелия, г. Петрозаводск, Россия;
Станкевич А.И., зав. отделом эпидемиологии Центра гигиены и эпидемиологии в Псковской области, г. Псков, Россия;
Калинина Е.Л., начальник отдела эпидемиологического надзора Управления Роспотребнадзора по Псковской области, г. Псков, Россия;
Викс Р., доктор наук, старший научный сотрудник Норвежского института общественного здравоохранения, Осло, Норвегия;
Andreassen A.K., доктор наук, профессор, старший научный сотрудник Норвежского института общественного здравоохранения, Осло, Норвегия.

Authors:

Tokarevich N.K., PhD, MD (Medicine), Professor, Head of the Laboratory of Zoonoses, St. Petersburg Pasteur Institute, St. Petersburg, Russian Federation;
Blinova O.V., PhD (Chemistry), Junior Researcher, Laboratory of Zoonoses, St. Petersburg Pasteur Institute, St. Petersburg, Russian Federation;
Stoyanova N.A., PhD (Medicine), Leading Researcher, Laboratory of Zoonoses, St. Petersburg Pasteur Institute, St. Petersburg, Russian Federation;
Baimova R.R., Junior Researcher, Laboratory of Zoonoses, St. Petersburg Pasteur Institute, St. Petersburg, Russian Federation;
Siuizumova E.A., Junior Researcher, Laboratory of Zoonoses, St. Petersburg Pasteur Institute, St. Petersburg, Russian Federation;
Lomonosova V.I., Junior Researcher, Laboratory of Zoonoses, St. Petersburg Pasteur Institute, St. Petersburg, Russian Federation;
Tronin A.A., PhD, MD (Geology and Mineralogy), Director, Scientific Research Center for Ecological Safety, Russian Academy of Sciences, St. Petersburg, Russian Federation;
Buzinov R.V., PhD, MD (Medicine), Associate Professor, Deputy Director for Development, Federal Service for Supervision of Consumer Rights Protection and Human Welfare in Arkhangelsk Region, Arkhangelsk, Russian Federation;
Sokolova O.V., Deputy Head of Epidemiological Surveillance Department, Federal Service for Supervision of Consumer Rights Protection and Human Welfare in Arkhangelsk Oblast, Arkhangelsk, Russian Federation;
Gnativ B.R., Deputy Head of Epidemiological Surveillance Department of Rospotrebnadzor Office for the Republic of Komi, Syktyvkar, Russian Federation;
Buts L.V., Head of Epidemiological Surveillance Department, Federal Service for Supervision of Consumer Rights Protection and Human Welfare in Leningrad Oblast, St. Petersburg, Russian Federation;
Bubnova L.A., Head of the Epidemiology Department, Center of Hygiene and Epidemiology in the Karelia Republic, Petrozavodsk, Russian Federation;
Safonova O.S., Biologist, Center of Hygiene and Epidemiology in the Karelia Republic, Petrozavodsk, Russian Federation;
Stankevich A.I., Head of Epidemiology Department, Center for Hygiene and Epidemiology in Pskov Oblast, Pskov, Russian Federation;
Kalinina E.L., Head of the Epidemiological Surveillance Department, Federal Service for Supervision of Consumer Rights Protection and Human Welfare in Pskov Oblast, Pskov, Russian Federation;
Vikse R., D.Sc., Senior Researcher, Norwegian Institute of Public Health, Oslo, Norway;
Andreassen A.K., D.Sc., Professor, Senior Researcher, Norwegian Institute of Public Health, Oslo, Norway.