

and their analysis were carried out using Microflex LT (Bruker) and its programs v. 3.3.64 and v. 3.3.65.

At the first stage we created 2 databases of mass spectra of reference strains: 1) saprophytes of the genus *Bacillus* and 2) strains of *B. anthracis*. When carrying out “blind” tests we revealed that fragments of peptide complexes over the range 2–12 000 Da in all representatives of both groups practically did not differ because of high degree of affinity. Thus, strains of closely related saprophytes were identified as *B. anthracis* and strains with high indicator SV on the contrary as saprophytes. When all spectra of cultures of both groups were pooled, identification became more correct, allowing to obtain the highest values of SV for strains of one species. The most optimum results of specific identification were obtained when identification of cultures was carried out using the program MALDI Biotyper RIC and construction of MsP-dendrogram was carried out using the program FlexAnalysis. In obtained dendrograms samples under study were clearly clustered with one of bacilli species represented in the base.

Thus, perfection of the scheme of reliable identification of *B. anthracis*, including accurate differentiation from closely related bacilli on the basis of MALDI TOF MS continues to remain urgent.

4.14 doi: 10.15789/2220-7619-2018-4-4.14

SEARCH FOR SPECIES-SPECIFIC MARKERS FOR *BACILLUS ANTHRACIS* BY MALDI-TOF MASS SPECTROMETRY

E.A. Koteneva, O.I. Tsygankova, A.V. Kalinin

Stavropol Plague Control Research Institute, Stavropol, Russia

The use of the sensitive and rapid method of MALDI-TOF MS for identification of cultures of the causative agent of anthrax requires not only strict specificity, but also universality for all strains irrespective of their intraspecific variability of phenotypic properties.

The aim of the work was to reveal species-specific signals, common to all *B. anthracis* strains with various complexes of phenotypic properties.

We used 37 strains which included strains atypical in capsule formation, toxin production, nutritional requirements, activities of protease, lecithinase and hemolysins, ability to hydrolyze carbohydrates, as well as strains with different MLVA- and SNP-genotypes. Samples were prepared by lysis of 18-hour cultures in 80% TFA followed by ultra-micro-centrifuge filtration. The studies were carried out using Microflex LT instrument (Bruker). Collection of mass spectra and analysis of data were carried out using the programs v. 3.3.64 and v. 3.3.65. Analysis of spectra for frequency of signals was carried out using the program Microbe MS.

The occurrence of various combinations of phenotypic properties made it possible to discriminate 11 phenotypes. Individual spectra of each of these phenotypes (20 spectra of each strain) were analyzed and peak frequency was determined. For the further analysis we used peaks occurring at the frequency $\geq 95\%$, with their numbers in various groups varying from 2 to 32.

When comparing the peak frequency of all the 11 phenotypic groups we revealed the absence of common peaks with the frequency $\geq 95\%$. The distribution of signals which were identified in all the groups most often were as follows: 2601 Da — 82.2%; 4367 Da — 81.7%; 4666 Da — 76.4%; 6445 Da — 73.8%; 5206 Da — 72.8%. Earlier these peaks were not considered as specific markers of *B. anthracis*. The approach to choose markers we used when analyzing strains with a great number of phenotypic groups, including

rare strains, may account for this. Markers of the system of ribosomal proteins, SASP and histone proteins, earlier described as species-specific markers, also occur in the spectra of strains from various groups, but at much lower frequency, and that may be connected with the production of various proteins or with various levels of their expression.

Thus, selection of species-specific peaks for identification of *B. anthracis* strains should be carried out taking into account the variability of their biological properties.

4.15 doi: 10.15789/2220-7619-2018-4-4.15

ETIOLOGICAL CHARACTERISTICS OF MALARIA AND PREVALENCE OF HEMOGLOBINOPATHIES IN PATIENTS IN THE REPUBLIC OF GUINEA

A.E. Levkovsky^{1,3}, D.A. Lioznov^{2,3}, A.H. Diallo¹, T.S. Sow¹, H.K. Diallo¹, V.G. Dedkov³

¹Hospital RUSAL FRIGUIA, Fria, Republic of Guinea;

²Smorodintsev Research Institute of Influenza; ³St. Petersburg Pasteur Institute, St. Petersburg, Russia

According to WHO, in 2016, malaria affected 216 million people in 91 countries, which is 5 million more than in 2015. The number of deaths from malaria in 2016 was 445 000 people. 90% of cases and 91% of deaths from malaria was from Africa.

There are more than 50 different types of hereditary hemoglobinopathies. They are most often found in regions with a tropical and subtropical climate, which correspond to geographic regions endemic for malaria.

The aim of our study was to determine the etiological structure of malaria and to assess the prevalence and variants of hemoglobinopathies in patients with malaria in the territory of the subprefecture Fria of the Republic of Guinea.

The study included 300 cases of malaria aged 0 to 70 years, from the hospital “RUSAL FRIGUIA” in town Fria from May to December 2017. Malaria was determined by a rapid test for the differentiated determination of antigen *P. falciparum* and pan-malarial antigen, with verification and validation of parasitemia by the method of thick drop and smear. The species belonging to the plasmodium was confirmed by the PCR method followed by sequencing. The type of hemoglobin was determined by method of electrophoresis.

The average age of patients was 15.8 years (from 1 month to 65 years), men — 53%. In 99% cases causative agent was *P. falciparum*, with parasitemia from 16 to 20 000 tr/ μ L. Hemoglobinopathy revealed in 20% of patients, first of all, sickle-cell anemia (85%). Lethal outcome was registered in 11 patients at the age from 2 to 14 years.

High parasitemia was associated with a more severe course of the disease. In patients with concomitant hemoglobinopathy revealed a less severe clinical course of malaria, characterized by relatively small parasitemia.

100% dominance of *P. falciparum* in patients with malaria in this region defines clinical vigilance regarding the severity of the course and the prognosis of the disease. Identifying concomitant hemoglobinopathies allow us to predict a favorable prognosis of malaria.

4.16 doi: 10.15789/2220-7619-2018-4-4.16

PECULIARITIES OF MASS SPECTROMETRIC ANALYSIS OF BRUCELLA S- AND L-FORMS

A.S. Ostyak, N.L. Barannikova, K.Yu. Yastremskaya, N.G. Gefan, N.A. Mikhailova, S.V. Balakhonov

Irkutsk Antiplague Research Institute, Irkutsk, Russia

The causative agent of brucellosis, like many bacteria, is able to transform from S- and R- forms into L-form under the influence of various factors changing its biological