

**EVALUATION OF SERUM LEVELS OF IL-6 AND ADIPONECTIN IN  
COVID-19 PATIENTS AND THEIR RELATIONSHIP WITH DISEASE  
SEVERITY**

Norouzian M.

Sharifi-Sarasiabi K.

Najafi-Asl M.

Hassani Azad M.

Estabraghnia Babaki H.

Hormozgan University of Medical Sciences, Bandar Abbas, Iran

**ОЦЕНКА СЫВОРОТОЧНЫХ УРОВНЕЙ ИЛ-6 И АДИПОНЕКТИНА  
У ПАЦИЕНТОВ С COVID-19 И ИХ СВЯЗИ С ТЯЖЕСТЬЮ  
ЗАБОЛЕВАНИЯ**

Норузьян М.

Шарифи-Сарасиаби К.

Наджафи-Асл М.

Хассани Азад М.

Эстабрагния Бабаки Х.

Гормозганский университет медицинских наук, Бендер-Аббас, Иран

## Abstract

**Background:** The SARS-CoV-2 pandemic has prompted researchers around the world to identify risk factors associated with disease severity and mortality. Results suggest that COVID-19 mortality might be due to a ‘cytokine storm’ involving IL-6, and that obesity can be considered a risk factor for COVID-19 prevalence, severity, and mortality. The current study aimed to evaluate the serum levels of IL-6 and adiponectin in patients and their relationship with disease progression. **Methods:** ELISA was used to assess the levels of IL-6 and adiponectin in serum samples from a control group and from patients with COVID-19 at the time of admission to ICU or non-ICU wards. The results were analyzed using the Mann-Whitney and Spearman tests. **Results:** Mean serum levels of adiponectin in patients admitted to ICU ( $10.18 \pm 15.4$  ng/ml) were significantly higher than patients admitted to non-ICU wards ( $3.14 \pm 3$  ng/ml,  $p = 0.001$ ). Mean serum IL-6 levels showed a similar pattern, however the difference was not statistically significant ( $p = 0.18$ ). In addition, a significant direct correlation was observed between adiponectin expression and IL-6 ( $R = 0.2$ ,  $p = 0.03$ ). **Conclusion:** The results of this study show that serum levels of adiponectin in COVID-19 patients with severe lung involvement were significantly higher than those with less lung involvement. This finding is of high importance mainly due to the critical role of the lungs in adiponectin signaling, and as a result, adiponectin disorders may be associated with pulmonary complications in COVID-19 patients.

**Keywords:** COVID-19, IL-6, adiponectin, intensive care unit, disease severity, lung involvement

**Резюме. История вопроса:** Пандемия SARS-CoV-2 инфекции побудила исследователей во всем мире выявить факторы риска, связанные с тяжестью заболевания и смертностью. Результаты показывают, что смертность от COVID-19 может быть связана с «цитокиновым штормом» с участием ИЛ-6, и что ожирение можно рассматривать как фактор риска

распространенности, тяжести и смертности от COVID-19. Настоящее исследование было направлено на оценку уровней ИЛ-6 и адипонектина в сыворотке у пациентов и их связь с прогрессированием COVID-19. Методы: ИФА использовался для оценки уровней ИЛ-6 и адипонектина в образцах сыворотки контрольной группы и пациентов с COVID-19 во время поступления в отделение интенсивной терапии или в обычные инфекционные отделения. Результаты анализировали с использованием тестов Манна-Уитни и Спирмена. **Результаты:** средние уровни адипонектина в сыворотке у пациентов, поступивших в отделение интенсивной терапии ( $10,18 \pm 15,4$  нг/мл), были значительно выше, чем у пациентов, поступивших в обычные инфекционные отделения ( $3,14 \pm 3$  нг/мл,  $P = 0,001$ ). Средние уровни ИЛ-6 в сыворотке были сравнимыми, без статистически достоверных различий ( $P = 0,18$ ). Кроме того, отмечена достоверная прямая корреляция между уровнем в сыворотке адипонектина и ИЛ-6 ( $R = 0,2$ ,  $P = 0,03$ ). **Вывод:** результаты настоящего исследования показали, что уровни адипонектина в сыворотке у пациентов с COVID-19 с тяжелым поражением легких были значительно выше, чем у пациентов с меньшей степенью поражения легких. Полученные данные имеют большое значение главным образом из-за критической роли легких в передаче сигналов от адипонектина, и, как следствие, нарушения в системе адипонектина могут быть связаны с легочными осложнениями у пациентов с COVID-19.

**Ключевые слова:** COVID-19, ИЛ-6, адипонектин, отделение реанимации, тяжесть заболевания, поражение легких.

## Introduction

COVID-19 is a recently discovered infectious disease caused by a new coronavirus called SARS-CoV-2 [1, 2]. In addition to older age, obesity has been revealed as one of the most important risk factors contributing to severe forms of COVID-19 [3]. In a retrospective study of 85 patients with COVID-19, obesity was reported as a risk factor for admission to the intensive care unit [4]. Although the mechanisms have yet to be elucidated, studies suggest that obesity can be considered a risk factor for acute respiratory failure [5], and that obesity might be associated with COVID-19 severity [6]. On the other hand, in people with lung injury, obesity has shown to be associated with better outcomes, a phenomenon called the ‘obesity paradox’ [7].

Adipose tissue is active in the processes of inflammation and immune response due to secretion of substances called ‘adipocytokines’ such as adiponectin, leptin, TNF- $\alpha$ , IL-6, complement factors, growth factors, and adhesive molecules [8]. Therefore, abnormal expression of adipocytokines secreted from adipose tissue may contribute to the ‘cytokine storm’ that characterizes severe COVID-19 forms. Several studies have shown the anti-inflammatory role of adiponectin in lung diseases, such as chronic obstructive pulmonary disease (COPD), emphysema, and lung cancer [9, 10]. The anti-inflammatory function of adiponectin has been shown in studies on lung cells *in vitro* [11, 12].

Mouse models with adiponectin deficiency also develop pulmonary dysfunction and systemic inflammation [13]. In fact, adiponectin has been shown to: decrease the proliferation of innate and acquired immune cells, as well as suppress the production of inflammatory mediators such as IL-6, IL-2, and TNF- $\alpha$ ; and to increase the production of immunosuppressive cytokines such as IL-10 [14, 15]. On the other hand, a significant association has also been reported between decreased blood levels of adiponectin and increased prevalence of cardiovascular disorders and obesity-related diabetes. Decreased

levels of adiponectin have also been associated with increased IL-6 levels [16-18].

However, recent studies suggest that adiponectin may also stimulate the production of inflammatory agents, and that elevated adiponectin levels might be associated with increased inflammation in inflammatory diseases such as autoimmune disorders. For example, elevated levels of adiponectin in serum and synovial fluid of patients with rheumatoid arthritis have shown to be directly related to serum CRP levels [19, 20]. Adiponectin has also been reported to increase the production of IL-6 in human synovial fibroblasts by activating the AMPK, p38MAPK, and NF- $\kappa$ B signaling pathways [21]. In type I diabetes, elevated levels of adiponectin have been associated with increased levels of IL-6 and CRP [22]. A direct relationship has also been described between elevated serum adiponectin levels and viral load and systemic inflammation in patients with chronic hepatitis infection [23, 24].

Since the interaction between adiponectin and inflammatory factors is not yet fully understood, further studies are needed especially in the context of COVID-19 infection causing pulmonary complications, such as pneumonia and, in the most severe cases, ARDS [25]. The results of a study involving 150 COVID-19 patients to determine the clinical predictors of disease mortality showed that in deceased patients, serum IL-6 levels were higher than in discharged patients; this implies that cytokine storm may be the leading cause of death in COVID-19 patients [26]. Therefore, several clinical trials have been conducted to find a useful treatment to reduce cytokine levels [27]. Only one study investigated the link between adiponectin and inflammatory cytokines in COVID-19 patients with acute respiratory distress. Blot et al. reported a negative association between adiponectin and IL-6 in COVID-19 patients [28]. Since reports on the relationship between inflammatory cytokines and adiponectin in COVID-19 patients are limited, this study aimed to evaluate the

serum levels of inflammatory cytokine IL-6 and adiponectin in patients with COVID-19 and to elucidate their relationship with disease progression.

## **Method and materials**

### **Study population**

In this study, serum samples from 80 patients with COVID-19, admitted to ICU or non-ICU wards of Shahid Mohammadi Hospital of Bandar Abbas, were collected on the first day of hospitalization. None of the patients received COVID-19 hospital treatment before sampling (use of medications by patients at home was however unavoidable). Serum samples were divided into appropriate tubes and kept at  $-70^{\circ}\text{C}$  until ELISA. Prior to sampling, a written informed consent was obtained from each patient. Definitive diagnosis of COVID-19 was based on positive PCR and lung involvement through chest CT scan. Control samples of healthy volunteers were also included.

All procedures performed in the current study were approved by the Ethics Committee of Hormozgan University of Medical Sciences (IR.HUMS.REC.1399.195) in accordance with the 1964 Helsinki declaration and its later amendments.

### **ELISA**

Serum levels of IL-6 and adiponectin were determined using commercial ELISA kits, based on the manufacturer's instructions (ZellBio, Germany). The sensitivities of the IL-6 and adiponectin assays were 30-960 pg/ml and 2-64 mg/l, respectively. Serum samples and standards were added to wells of plates containing antibodies against IL-6 or adiponectin. After incubation and washing, HRP-conjugated antibody was added to the plates. Following incubation and washing, substrate was then added. In the next step, enzyme inhibitor solution was added, and plates were read by ELISA reader at a wavelength of 450 nm.

### **Statistical analysis**

Data were analyzed using the Mann-Whitney and Spearman non-parametric statistical tests with SPSS software version 16. Significance was designated as  $p < 0.05$ .

## Results

The study population involved: 80 patients (40 admitted to the ICU, and 40 admitted to non-ICU wards); and 5 healthy controls. All groups were matched according to demographic variables, such as age and sex. A total of 41 men and 44 women were studied. The mean age of the study population was  $28.45 \pm 5.83$  years. The mean ages were:  $60.2 \pm 1.7$  in the ICU group;  $56.8 \pm 1.7$  in the non-ICU patients; and  $44.2 \pm 18.3$  years in the control group. There was no difference between groups according to different variables (Table 1). Detailed information on the study population is presented in Table 1. However, due to a lack of information on patient weights and heights, we were not able to compare groups in terms of body mass index (BMI).

### Serum IL-6 levels

The means ( $\pm$  standard deviation) of IL-6 level in ICU and non-ICU patients were  $48.7 \pm 2.7$  and  $41.9 \pm 2.7$  pg/ml, respectively, which was not statistically significant nor in control subjects (Fig. 1) ( $p = 0.18$ ). IL-6 expression in young people (less than 45 years) admitted to the ICU was significantly higher than young people admitted to the non-ICU wards ( $p = 0.02$ ). In ICU hospitalized patients with diabetes and hypertension, IL-6 levels were significantly higher compared to patients without underlying disease ( $p = 0.01$ ,  $p = 0.02$ , respectively).

### Serum adiponectin levels

ICU and non-ICU hospitalized patients had higher adiponectin levels compared to the control group ( $p < 0.001$  and  $p = 0.002$ , respectively). Moreover, the mean serum level of adiponectin in ICU patients was significantly higher than in non-ICU patients ( $10.4 \pm 15.4$  and  $3.1 \pm 3$  mg/l, respectively,  $p = 0.001$ ) (Fig. 1), and it was significant in both age groups

(younger  $p = 0.003$ , older  $p = 0.03$ ). However, significant differences in adiponectin level in patients with comorbidities, compared to those without comorbidities, were not seen. The results of our study show that there was a significant positive relationship between IL-6 and adiponectin in COVID-19 patients ( $p = 0.03$ ,  $R = 0.232$ ) (Fig. 2).

## Discussion

Due to mortality (5-7%) and high ICU admission (9-11%) among patients infected with SARS-CoV-2 [29], all necessary measures should be taken to control the COVID-19 pandemic. Since COVID-19 can cause mild illness, or a very acute respiratory syndrome leading to release of proinflammatory cytokines including IL-6 and TNF- $\alpha$  [26], and considering obesity as a potential risk factor for developing serious illness with SARS-CoV-2 infection [4], a possible mechanism through abnormal secretion of adipokines via adipose tissue can be proposed. Therefore, in this study, serum levels of IL-6 and adiponectin (the most plentiful adipokine released by adipose tissue) in COVID-19 patients, and their relationship with disease severity, were evaluated. Our study showed that hospitalized COVID-19 patients had significantly elevated serum adiponectin levels compared to healthy controls. Moreover, adiponectin levels were significantly higher in ICU patients compared to patients admitted to non-ICU wards; this implies a possible relationship between adiponectin and disease severity.

Serum IL-6 levels were also higher in the ICU group compared to the non-ICU group. The difference, however, was not statistically significant. It should be noted that the small size of our study population (80 individuals), along with self-medication at home and arbitrary consumption of anti-inflammatory drugs (such as nonsteroidal anti-inflammatory drugs, NSAIDs) likely influenced IL-6 expression. NSAIDs are well known as immunoregulators, but their mechanisms of action have not been fully determined. Previous studies have shown that NSAIDs down-regulate IL-6



production at the mRNA and protein levels in human PBMCs, T-cell clones, and chondrocytes [29, 30]. Moreover, in a mouse model of SARS-CoV-2 infection, it was shown that NSAID treatment decreased the production of a subset of inflammatory cytokines upregulated by infection, including IL-6 [31].

Adiponectin is an adipokine secreted in large quantities, and almost exclusively, from adipose tissue. As mentioned, protective functions of adiponectin have been shown in experimental disease models, such as atherosclerosis, diabetes, and inflammation [8]. Animal and *in vitro* studies have shown that adiponectin inhibits the production of proinflammatory cytokines, including TNF- $\alpha$  and IL-6 [14, 15, 30], while increasing anti-inflammatory mediators such as IL-10 [31]. Despite these observations, several studies have shown that adiponectin exerts proinflammatory effects under certain conditions [32, 33]. Several clinical studies have proposed a possible relationship between adiponectin levels and mortality rates in patients with acute dyspnea and cardiovascular diseases [34, 35].

In a large cross-sectional study of patients with pneumonia or sepsis admitted to intensive care, Waki et al. found that higher concentrations of systemic adiponectin in the early days of the disease were associated with low survival rates [36]. It has been proposed that higher concentrations of adiponectin on the first day of illness might indicate an inefficient response to disease induced stress, suggesting that an initial drop in adiponectin levels and subsequent return to baseline could be beneficial for patient survival. According to this hypothesis, the deceased patients in this study showed a slight increase in adiponectin levels between day 1 and day 6 of hospitalization [36]. Similarly, in the present study, the adiponectin levels in patients admitted to intensive care on the first day of hospitalization were higher compared to patients admitted to non-ICU wards. However, adiponectin might exert different effects on critical illnesses in different time periods. Therefore, further longitudinal studies are needed to evaluate the concentrations of adiponectin in different time periods

during the course of disease to determine its function in disease progression.

On the other hand, adiponectin has been shown to exhibit pro-inflammatory and anti-inflammatory activities [37]. Therefore, it could exert different effects under different inflammatory conditions. Patients with chronic inflammatory diseases, such as rheumatoid arthritis, SLE, type I diabetes, and inflammatory bowel disease, had elevated adiponectin levels compared with patients with metabolic disorders [20, 38-40]. These conditions point to a clear inconsistency, which is characterized by the presence of large quantities of adiponectin in an inflammatory context. In these patients, adiponectin levels are positively correlated to inflammatory markers. In addition, pro-inflammatory functions of adiponectin have been reported in the synovium of the joints and in the epithelium of the large intestine.

Adiponectin has been shown to have pro- and anti-inflammatory properties, however the effect of total adiponectin in chronic inflammatory diseases, such as autoimmune disorders, is controversial. It is worth mentioning that adiponectin consists of three isoforms: low molecular weight (trimers); intermediate molecular weight (hexamers); and high molecular weight (HMW or multimers) [41]. These isoforms may differ in their biological activities. For example, the HMW isoform is the most biologically active form in the regulation of insulin resistance [42]. For this reason, further studies are needed to investigate the possible roles of adiponectin isoforms in inflammatory responses.

Adiponectin and its receptors are expressed in different lung cells [43, 44]. Moreover, pulmonary endothelial cells contribute to transfer of adiponectin from the circulation into the lungs by expressing the cadherin molecule [45]. Therefore, the lung could be a target organ for adiponectin signaling, and as a result, adiponectin disorders might be associated with pulmonary complications in COVID-19 patients. However, further studies are needed to determine the role and regulation of adiponectin in inflammation and COVID-19 patients.

Investigation of adiponectin function and regulation, along with its possible pleiotropy, would help develop new methodologies for diagnosis and management of disease. The limitations of the present study include: a lack of BMI data; the small sizes of the patient and healthy control groups; and no evaluation of adiponectin isoforms. These limitations should be considered in future studies.

Altogether, the present study provides valuable data on the associations between adiponectin level, IL-6 level, and disease severity in COVID-19 patients with acute respiratory distress. However, the relationship between adiponectin and proinflammatory agents is still unclear. In addition, the precise anti-inflammatory or proinflammatory functions of systemic and local adiponectin, and their association with IL-6 expression, need to be elucidated in various diseases including COVID-19.

### **Acknowledgement**

The authors acknowledge the efforts of staff at hospitals affiliated with the Hormozgan University of Medical Sciences and also the patient participants in the study. This research was funded by the Hormozgan University of Medical Sciences (IR.HUMS.REC.1399.195).

### **Conflict of interest**

The authors declare that there are no conflicts of interest.

РИСУНКИ

Figure 1





Figure 2



### ПОДПИСИ К РИСУНКАМ

Figure 1. Comparison of serum IL-6 and adiponectin levels in COVID-19 patients admitted to ICU wards, non-ICU wards, and control group. Eighty patients were divided into two groups: ICU hospitalized (n = 40) and non-ICU (n = 40). IL-6 (A) and adiponectin (B) levels were compared between the two groups. The values in the figure are expressed as Mean  $\pm$  SEM. \*\*\* *p* value < 0.001.

Figure 2. Spearman's correlation between serum levels of IL-6 and adiponectin in COVID-19 patients (n=80).

## ТАБЛИЦЫ

Table 1: Demographics and baseline characteristics of COVID-19 patients

	All patients (n=80)	ICU care (n=40)	Non-ICU care (n=40)	Healthy controls (n=5)	<i>p</i> value
<b>Characteristics</b>					
Age	58.5±1.7 (24-94)	60.2±1.7 (26-88)	56.8±1.7(24-94)	44.2±18.3(27-75)	0.33
Sex					
Male	39 (48.8%)	20(50%)	19 (47.5%)	2 (40%)	-
Female	41 (51.2%)	20 (50%)	21 (52.5%)	3 (60%)	-
Diabetes	30 (37.5%)	14 (35%)	16 (40%)	-	0.64
Cardiovascular disease	26 (32.5%)	12 (30%)	14 (35%)	-	0.63
Hypertension	32 (40%)	19 (47.5%)	13 (32.5%)	-	0.17
<b>Signs and symptoms</b>					
Fever	40 (50%)	17 (42.5%)	23 (57.5%)	-	0.18
Cough	42 (52.5%)	20 (50%)	22 (55%)	-	0.65
Dyspnea	1 (1.2%)	0	1 (2.5%)	-	0.31
Respiratory distress	62 (77%)	32 (80%)	30 (75%)	-	0.59
Myalgia	20 (25%)	11 (27.5%)	9 (22.5%)	-	0.60
Headache	9 (11.2%)	5 (12.5%)	4 (10%)	-	0.72
Vertigo	8 (10%)	5 (12.5%)	3 (7.5%)	-	0.45
Fatigue	14 (17.5%)	7 (17.5%)	7 (17.5%)	-	1.00
Anorexia	11 (13.8%)	5 (12.5%)	6 (15%)	-	0.74
Loss of taste	1 (1.2%)	0	1 (2.5%)	-	0.31
Diarrhea	9 (11.2%)	3 (7.5%)	6 (15%)	-	0.29
Vomiting	4 (5%)	1 (2.5%)	3 (7.5%)	-	0.30
Chest pain	5 (6.2%)	2 (5%)	3 (7.5%)	-	0.64
Death	39 (48.8%)	32 (80%)	7 (17.5%)	-	<0.0001
IL-6 (pg/ml)	45.3±2.7 (3-155.5)	48.7±2.7 (10-155.5)	41.9±2.7 (3-135)		0.18
Adiponectin (µg/ml)	6.7±11.6 (0.0-84.8)	10.18±15.4 (0.8-84.8)	3.14±3 (0.0-14.5)		0.001
	Data are mean±SD (standard deviation), with n (%), or n/N (%), where N is the total number of patients with available data. <i>p</i> values comparing ICU care and non-ICU care are from Mann-Whitney U test. COVID-19 – coronavirus disease 2019. ICU – intensive care unit.				

## ТИТУЛЬНЫЙ ЛИСТ\_МЕТАДААННЫЕ

Evaluation of serum levels of IL-6 and adiponectin in COVID-19 patients and their relationship with disease severity

### **Блок 1.** Информация об авторе, ответственном за переписку

Marzieh Norouzian, Ph.D., Assistant Professor of Medical Immunology

Department of Laboratory Sciences, School of Allied Medical Sciences,  
Hormozgan University of Medical Sciences, Bandar Abbas, Iran., P.O. Box:  
7919693116, Bandar Abbas, Iran. Tel: +98 763366367; Fax: +98763366367

E-mail address: marzieh.norouzi@gmail.com

### **Блок 2.** Информация об авторах

Marzieh Norouzian, Department of Laboratory Sciences, School of Allied  
Medical Sciences, Hormozgan University of Medical Sciences, Bandar Abbas,  
Iran

Khojasteh Sharifi-Sarasiabi, Infectious & Tropical Diseases Research Center,  
Hormozgan University of Medical Sciences, Bandar Abbas, Iran

Majid Najafi-Asl, Department of Laboratory Sciences, School of Allied  
Medical Sciences, Hormozgan University of Medical Sciences, Bandar Abbas,  
Iran; Infectious & Tropical Diseases Research Center, Hormozgan University of  
Medical Sciences, Bandar Abbas, Iran

Mehdi Hassani Azad, Infectious & Tropical Diseases Research Center,  
Hormozgan University of Medical Sciences, Bandar Abbas, Iran

Hamideh Estabraghnia Babaki, Department of Internal Medicine, Shahid  
Mohammadi Hospital, Faculty of Medicine, Hormozgan University of Medical  
Sciences, Bandar Abbas, Iran

### **Блок 3.** Метаданные статьи

Оригинальная статья

**30.07.2021**



## REFERENCES

Reference sequence number	Title of a publication and source in English	Reference's URL
Ajuwon, K. M. and M. E. Spurlock. Adiponectin inhibits LPS-induced NF- $\kappa$ B activation and IL-6 production and increases PPAR $\gamma$ 2 expression in adipocytes. <i>American Journal of Physiology-Regulatory, Integrative and Comparative Physiology</i> , 2005, vol. 288, no 5, pp R1220-R1225.	-	<a href="https://journals.physiology.org/doi/full/10.1152/ajpregu.00397.2004">https://journals.physiology.org/doi/full/10.1152/ajpregu.00397.2004</a>
Baud, D., X. Qi, K. Nielsen-Saines, D. Musso, L. Pomar and G. Favre. Real estimates of mortality following COVID-19 infection. <i>The Lancet infectious diseases</i> , 2020, vol. 20, no 7, pp 773.	-	<a href="https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(20)30195-X/fulltext">https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(20)30195-X/fulltext</a>
Bianco, A., G. Mazzeola, V. Turchiarelli, E. Nigro, G. Corbi, O. Scudiero, M. Sofia and A. Daniele. Adiponectin: an attractive marker for metabolic disorders in Chronic Obstructive Pulmonary Disease (COPD). <i>Nutrients</i> , 2013, vol. 5, no 10, pp 4115-4125.	-	<a href="https://www.mdpi.com/58942">https://www.mdpi.com/58942</a>
Blot, M., M. David, M. Nguyen, A. Bourredjem, C. Binquet and L. Piroth. Are adipokines the missing link between obesity, immune response, and outcomes in severe COVID-19? <i>Int J Obes (Lond)</i> , 2021, vol. 45, no 9, pp 2126-2131.	-	<a href="https://www.nature.com/articles/s41366-021-00868-5">https://www.nature.com/articles/s41366-021-00868-5</a>
Chen, J. S., M. M. Alfajaro, R. D. Chow, J. Wei, R. B. Filler, S. C. Eisenbarth and C. B. Wilen. Non-steroidal anti-inflammatory drugs dampen the cytokine and antibody response to SARS-CoV-2 infection. <i>J Virol</i> , 2021, vol. 95, no 7.	-	<a href="https://journals.asm.org/doi/full/10.1128/JVI.00014-21">https://journals.asm.org/doi/full/10.1128/JVI.00014-21</a>
Chiang, C. H., J. S. Lai, S. H. Hung, L. T. Lee, J. C. Sheu and K. C. Huang. Serum adiponectin levels are associated with hepatitis B viral load in overweight to obese hepatitis B virus carriers. <i>Obesity</i> , 2013, vol. 21, no 2, pp 291-296.	-	<a href="https://onlinelibrary.wiley.com/doi/full/10.1002/oby.20000">https://onlinelibrary.wiley.com/doi/full/10.1002/oby.20000</a>
Choi, H. M., H. M. Doss and K. S. Kim. Multifaceted Physiological Roles of Adiponectin in Inflammation and Diseases. <i>Int J Mol Sci</i> , 2020, vol. 21, no 4.	-	<a href="https://www.mdpi.com/638980">https://www.mdpi.com/638980</a>
Daniele, A., A. De Rosa, E. Nigro, O. Scudiero, M. Capasso, M. Masullo, G. De Laurentiis, G. Oriani, M. Sofia and A. Bianco. Adiponectin	-	<a href="https://pubmed.ncbi.nlm.nih.gov/2223">https://pubmed.ncbi.nlm.nih.gov/2223</a>

oligomerization state and adiponectin receptors airway expression in chronic obstructive pulmonary disease. The international journal of biochemistry & cell biology,2012, vol. 44, no 3, pp 563-569.		3975/
Dieplinger, B., A. Gegenhuber, G. Kaar, W. Poelz, M. Haltmayer and T. Mueller. Prognostic value of established and novel biomarkers in patients with shortness of breath attending an emergency department. Clin Biochem,2010, vol. 43, no 9, pp 714-719.	-	<a href="https://www.sciencedirect.com/science/article/abs/pii/S0009912010000615">https://www.sciencedirect.com/science/article/abs/pii/S0009912010000615</a>
Ehling, A., A. Schäffler, H. Herfarth, I. H. Tarner, S. Anders, O. Distler, G. Paul, J. Distler, S. Gay and J. Schölmerich. The potential of adiponectin in driving arthritis. The Journal of Immunology,2006, vol. 176, no 7, pp 4468-4478.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/16547285/">https://pubmed.ncbi.nlm.nih.gov/16547285/</a>
Furuta, M., M. Tamai, T. Hanabusa, Y. Yamamoto, K. Nanjo and T. Sanke. Serum adiponectin is associated with fasting serum C-peptide in non-obese diabetic patients. Diabetes research and clinical practice,2006, vol. 72, no 3, pp 302-307.	-	<a href="https://www.sciencedirect.com/science/article/abs/pii/S0168822705004316">https://www.sciencedirect.com/science/article/abs/pii/S0168822705004316</a>
Guan, W.-j., Z.-y. Ni, Y. Hu, W.-h. Liang, C.-q. Ou, J.-x. He, L. Liu, H. Shan, C.-l. Lei and D. S. Hui. Clinical characteristics of coronavirus disease 2019 in China. New England Journal of Medicine,2020, vol. 382, no 18, pp 1708-1720.	-	<a href="https://www.nejm.org/doi/full/10.1056/nejmoa2002032">https://www.nejm.org/doi/full/10.1056/nejmoa2002032</a>
Hadjadj, S., R. Aubert, F. Fumeron, F. Pean, J. Tichet, R. Roussel and M. Marre. Increased plasma adiponectin concentrations are associated with microangiopathy in type 1 diabetic subjects. Diabetologia,2005, vol. 48, no 6, pp 1088-1092.	-	<a href="https://link.springer.com/article/10.1007/s00125-005-1747-x">https://link.springer.com/article/10.1007/s00125-005-1747-x</a>
Huang, C., Y. Wang, X. Li, L. Ren, J. Zhao, Y. Hu, L. Zhang, G. Fan, J. Xu and X. Gu. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. The Lancet,2020, vol. 395, no 10223, pp 497-506.	-	<a href="https://www.thelancet.com/journals/lanet/article/PIIS0140-6736(20)30183-5/fulltext">https://www.thelancet.com/journals/lanet/article/PIIS0140-6736(20)30183-5/fulltext</a>
Hug, C., J. Wang, N. S. Ahmad, J. S. Bogan, T.-S. Tsao and H. F. Lodish. T-cadherin is a receptor for hexameric and high-molecular-weight forms of Acrp30/adiponectin. Proceedings of the National Academy of Sciences,2004, vol. 101, no 28, pp 10308-10313.	-	<a href="https://www.pnas.org/doi/full/10.1073/pnas.0403382101">https://www.pnas.org/doi/full/10.1073/pnas.0403382101</a>

Jordan, R. E., P. Adab and K. Cheng. Covid-19: risk factors for severe disease and death, British Medical Journal Publishing Group, 2020.	-	<a href="https://www.bmj.com/content/368/bmj.m1198">https://www.bmj.com/content/368/bmj.m1198</a>
Krakoff, J., T. Funahashi, C. D. Stehouwer, C. G. Schalkwijk, S. Tanaka, Y. Matsuzawa, S. Kobes, P. A. Tataranni, R. L. Hanson and W. C. Knowler. Inflammatory markers, adiponectin, and risk of type 2 diabetes in the Pima Indian. Diabetes care,2003, vol. 26, no 6, pp 1745-1751.	-	<a href="https://diabetesjournals.org/care/article/26/6/1745/26354/Inflammatory-Markers-Adiponectin-and-Risk-of-Type">https://diabetesjournals.org/care/article/26/6/1745/26354/Inflammatory-Markers-Adiponectin-and-Risk-of-Type</a>
Malhotra, A. and D. Hillman. Obesity and the lung: 3. Obesity, respiration and intensive care. Thorax,2008, vol. 63, no 10, pp 925-931.	-	<a href="https://thorax.bmj.com/content/63/10/925">https://thorax.bmj.com/content/63/10/925</a>
Mantzoros, C. S., T. Li, J. E. Manson, J. B. Meigs and F. B. Hu. Circulating adiponectin levels are associated with better glycemic control, more favorable lipid profile, and reduced inflammation in women with type 2 diabetes. The Journal of Clinical Endocrinology & Metabolism,2005, vol. 90, no 8, pp 4542-4548.	-	<a href="https://academic.oup.com/jcem/article/90/8/4542/3058890">https://academic.oup.com/jcem/article/90/8/4542/3058890</a>
Masaki, T., S. Chiba, H. Tatsukawa, T. Yasuda, H. Noguchi, M. Seike and H. Yoshimatsu. Adiponectin protects LPS-induced liver injury through modulation of TNF- $\alpha$ in KK-Ay obese mice. Hepatology,2004, vol. 40, no 1, pp 177-184.	-	<a href="https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/hep.20282">https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/hep.20282</a>
Mehta, P., D. F. McAuley, M. Brown, E. Sanchez, R. S. Tattersall and J. J. Manson. COVID-19: consider cytokine storm syndromes and immunosuppression. The Lancet,2020, vol. 395, no 10229, pp 1033-1034.	-	<a href="https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)30628-0/fulltext">https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)30628-0/fulltext</a>
Nigro, E., O. Scudiero, D. Sarnataro, G. Mazzearella, M. Sofia, A. Bianco and A. Daniele. Adiponectin affects lung epithelial A549 cell viability counteracting TNF $\alpha$ and IL-1 $\beta$ toxicity through AdipoR1. The international journal of biochemistry & cell biology,2013, vol. 45, no 6, pp 1145-1153.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/23500159/">https://pubmed.ncbi.nlm.nih.gov/23500159/</a>
Ohashi, K., N. Ouchi and Y. Matsuzawa. Anti-inflammatory and anti-atherogenic properties of adiponectin. Biochimie,2012, vol. 94, no 10, pp	-	<a href="https://pubmed.ncbi.nlm.nih.gov/2271">https://pubmed.ncbi.nlm.nih.gov/2271</a>

2137-2142.		3764/
Ohashi, K., R. Shibata, T. Murohara and N. Ouchi. Role of anti-inflammatory adipokines in obesity-related diseases. Trends Endocrinol Metab,2014, vol. 25, no 7, pp 348-355.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/24746980/">https://pubmed.ncbi.nlm.nih.gov/24746980/</a>
Otero, M., R. Lago, R. Gomez, F. Lago, C. Dieguez, J. J. Gomez-Reino and O. Gualillo. Changes in fat-derived hormones plasma concentrations: adiponectin, leptin, resistin, and visfatin in rheumatoid arthritis subjects. Annals of the Rheumatic Diseases,2006, vol. 65, no. 9, pp 1198-201.	-	<a href="https://ard.bmj.com/content/65/9/1198">https://ard.bmj.com/content/65/9/1198</a>
Pajvani, U. B., M. Hawkins, T. P. Combs, M. W. Rajala, T. Doebber, J. P. Berger, J. A. Wagner, M. Wu, A. Knopps and A. H. Xiang. Complex distribution, not absolute amount of adiponectin, correlates with thiazolidinedione-mediated improvement in insulin sensitivity. Journal of Biological Chemistry,2004, vol. 279, no 13, pp 12152-12162.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/14699128/">https://pubmed.ncbi.nlm.nih.gov/14699128/</a>
Pemmari, A., L. Tuure, M. Hämäläinen, T. Leppänen, K. Vuolteenaho, T. Moilanen and E. Moilanen. Comprehensive effects of ibuprofen on gene expression in chondrocytes as determined by RNA-Seq. Osteoarthritis and Cartilage,2019, vol. 27, pp S378.	-	<a href="https://rmdopen.bmj.com/content/7/3/e001657">https://rmdopen.bmj.com/content/7/3/e001657</a>
Petrakis, D., D. Margină, K. Tsarouhas, F. Tekos, M. Stan, D. Nikitovic, D. Kouretas, D. A. Spandidos and A. Tsatsakis. Obesity-a risk factor for increased COVID-19 prevalence, severity and lethality. Molecular Medicine Reports,2020, vol. 22, no. 1. Pp 9-19.	-	<a href="https://www.spandidos-publications.com/10.3892/mmr.2020.11127">https://www.spandidos-publications.com/10.3892/mmr.2020.11127</a>
Polito, R., E. Nigro, A. Elce, M. L. Monaco, P. Iacotucci, V. Carnovale, M. Comegna, M. Gelzo, F. Zarrilli and G. Corso. Adiponectin Expression Is Modulated by Long-Term Physical Activity in Adult Patients Affected by Cystic Fibrosis. Mediators of inflammation,2019, vol. 2019, pp 2153934.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/31582896/">https://pubmed.ncbi.nlm.nih.gov/31582896/</a>
Ruan, Q., K. Yang, W. Wang, L. Jiang and J. Song. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. Intensive care medicine,2020, pp 1-3.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/32125452/">https://pubmed.ncbi.nlm.nih.gov/32125452/</a>
Sada, K.-E., Y. Yamasaki, M. Maruyama, H. Sugiyama, M. Yamamura, Y. Maeshima and H. Makino. Altered levels of adipocytokines in association with insulin resistance in patients with systemic lupus erythematosus. The	-	<a href="https://pubmed.ncbi.nlm.nih.gov/16881111/">https://pubmed.ncbi.nlm.nih.gov/16881111/</a>

Journal of rheumatology,2006, vol. 33, no 8, pp 1545-1552.		
Schäffler, A., A. Ehling, E. Neumann, H. Herfarth, I. Tarner, J. Schölmerich, U. Müller-Ladner and S. Gay. Adipocytokines in Synovial FluidRESEARCH LETTER. Jama,2003, vol. 290, no 13, pp 1709-1710.	-	<a href="https://jamanetwork.com/journals/jama/article-abstract/197371">https://jamanetwork.com/journals/jama/article-abstract/197371</a>
Schnabel, R., C. M. Messow, E. Lubos, C. Espinola-Klein, H. J. Rupprecht, C. Bickel, C. Sinning, S. Tzikas, T. Keller, S. Genth-Zotz, K. J. Lackner, T. F. Münzel and S. Blankenberg. Association of adiponectin with adverse outcome in coronary artery disease patients: results from the AtheroGene study. Eur Heart J,2008, vol. 29, no 5, pp 649-657.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/18263867/">https://pubmed.ncbi.nlm.nih.gov/18263867/</a>
Simonnet, A., M. Chetboun, J. Poissy, V. Raverdy, J. Noulette, A. Duhamel, J. Labreuche, D. Mathieu, F. Pattou and M. Jourdain. High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. Obesity,2020, vol. 28, no. 7. pp 1195-1199.	-	<a href="https://onlinelibrary.wiley.com/doi/10.1002/oby.22831">https://onlinelibrary.wiley.com/doi/10.1002/oby.22831</a>
Sood, A., E. Dominic, C. Qualls, M. W. Steffes, B. Thyagarajan, L. J. Smith, C. E. Lewis and D. R. Jacobs Jr. Serum adiponectin is associated with adverse outcomes of asthma in men but not in women. Frontiers in pharmacology,2011, vol. 2, pp 55.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/22007173/">https://pubmed.ncbi.nlm.nih.gov/22007173/</a>
Takemura, Y., N. Ouchi, R. Shibata, T. Aprahamian, M. T. Kirber, R. S. Summer, S. Kihara and K. Walsh. Adiponectin modulates inflammatory reactions via calreticulin receptor–dependent clearance of early apoptotic bodies. The Journal of clinical investigation,2007, vol. 117, no 2, pp 375-386.	-	<a href="https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1770947/">https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1770947/</a>
Tang, C.-H., Y.-C. Chiu, T.-W. Tan, R.-S. Yang and W.-M. Fu. Adiponectin enhances IL-6 production in human synovial fibroblast via an AdipoR1 receptor, AMPK, p38, and NF-κB pathway. The Journal of Immunology,2007, vol. 179, no 8, pp 5483-5492.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/17911635/">https://pubmed.ncbi.nlm.nih.gov/17911635/</a>
Tiftikci, A., O. Atug, Y. Yilmaz, F. Eren, F. T. Ozdemir, S. Yapali, O. Ozdogan, C. A. Celikel, N. Imeryuz and N. Tozun. Serum levels of adipokines in patients with chronic HCV infection: relationship with steatosis and fibrosis. Archives of medical research,2009, vol. 40, no 4, pp 294-298.	-	<a href="https://bmcgastroenterol.biomedcentral.com/articles/10.1186/1471-230X-14-27">https://bmcgastroenterol.biomedcentral.com/articles/10.1186/1471-230X-14-27</a>

Tsuboi, I., H. Tanaka, M. Nakao, S. Shichijo and K. Itoh. Nonsteroidal anti-inflammatory drugs differentially regulate cytokine production in human lymphocytes: up-regulation of TNF, IFN-gamma and IL-2, in contrast to down-regulation of IL-6 production. <i>Cytokine</i> ,1995, vol. 7, no 4, pp 372-379.	-	<a href="https://api.semanticscholar.org/CorpusID:24976365">https://api.semanticscholar.org/CorpusID:24976365</a>
Vaduganathan, M., O. Vardeny, T. Michel, J. J. McMurray, M. A. Pfeffer and S. D. Solomon. Renin–angiotensin–aldosterone system inhibitors in patients with Covid-19. <i>New England Journal of Medicine</i> ,2020, vol. 382, no 17, pp 1653-1659.	-	<a href="https://www.nejm.org/doi/full/10.1056/nejmsr2005760">https://www.nejm.org/doi/full/10.1056/nejmsr2005760</a>
Walkey, A. J., T. W. Rice, J. Konter, N. Ouchi, R. Shibata, K. Walsh, B. P. deBoisblanc and R. Summer. Plasma adiponectin and mortality in critically ill subjects with acute respiratory failure. <i>Crit Care Med</i> ,2010, vol. 38, no 12, pp 2329-2334.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/20890191/">https://pubmed.ncbi.nlm.nih.gov/20890191/</a>
Wang, Y., K. S. Lam, M.-h. Yau and A. Xu. Post-translational modifications of adiponectin: mechanisms and functional implications. <i>Biochemical Journal</i> ,2008, vol. 409, no 3, pp 623-633.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/18177270/">https://pubmed.ncbi.nlm.nih.gov/18177270/</a>
Wolf, A. M., D. Wolf, H. Rumpold, B. Enrich and H. Tilg. Adiponectin induces the anti-inflammatory cytokines IL-10 and IL-1RA in human leukocytes. <i>Biochemical and biophysical research communications</i> ,2004, vol. 323, no 2, pp 630-635.	-	<a href="https://europepmc.org/article/med/15369797">https://europepmc.org/article/med/15369797</a>
Wozniak, S. E., L. L. Gee, M. S. Wachtel and E. E. Frezza. Adipose tissue: the new endocrine organ? A review article. <i>Dig Dis Sci</i> ,2009, vol. 54, no 9, pp 1847-1856.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/19052866/">https://pubmed.ncbi.nlm.nih.gov/19052866/</a>
Wulster-Radcliffe, M. C., K. M. Ajuwon, J. Wang, J. A. Christian and M. E. Spurlock. Adiponectin differentially regulates cytokines in porcine macrophages. <i>Biochemical and biophysical research communications</i> ,2004, vol. 316, no 3, pp 924-929.	-	<a href="https://europepmc.org/article/med/15033490">https://europepmc.org/article/med/15033490</a>
Yamamoto, K., T. Kiyohara, Y. Murayama, S. Kihara, Y. Okamoto, T. Funahashi, T. Ito, R. Nezu, S. Tsutsui and J. Miyagawa. Production of adiponectin, an anti-inflammatory protein, in mesenteric adipose tissue in Crohn’s disease. <i>Gut</i> ,2005, vol. 54, no 6, pp 789-796.	-	<a href="https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1774527/">https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1774527/</a>
Yamauchi, T., Y. Nio, T. Maki, M. Kobayashi, T. Takazawa, M. Iwabu, M. Okada-Iwabu, S. Kawamoto, N. Kubota and T. Kubota. Targeted disruption	-	<a href="https://www.nature.com/articles/nm15">https://www.nature.com/articles/nm15</a>

of AdipoR1 and AdipoR2 causes abrogation of adiponectin binding and metabolic actions. Nature medicine,2007, vol. 13, no 3, pp 332-339.		57
Zhi, G., W. Xin, W. Ying, X. Guohong and L. Shuying. "Obesity Paradox" in Acute Respiratory Distress Syndrome: Asystematic Review and Meta-Analysis. PLoS One,2016, vol. 11, no 9, pp e0163677.	-	<a href="https://pubmed.ncbi.nlm.nih.gov/27684705/">https://pubmed.ncbi.nlm.nih.gov/27684705/</a>